

Homogenization of Elliptic Boundary Value Problems in Lipschitz Domains

Carlos E. Kenig*

Zhongwei Shen†

Dedicated to the Memory of Björn Dahlberg

Abstract

In this paper we study the L^p boundary value problems for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$, where $\mathcal{L} = -\operatorname{div}(A\nabla)$ is a second order elliptic operator with real and symmetric coefficients. Assume that A is *periodic* in x_{d+1} and satisfies some minimal smoothness condition in the x_{d+1} variable, we show that the L^p Neumann and regularity problems are uniquely solvable for $1 < p < 2 + \delta$. We also present a new proof of Dahlberg's theorem on the L^p Dirichlet problem for $2 - \delta < p < \infty$ (Dahlberg's original unpublished proof is given in the Appendix). As the periodic and smoothness conditions are imposed only on the x_{d+1} variable, these results extend directly from $\mathbb{R}_+^{d+1}$ to regions above Lipschitz graphs. Consequently, by localization techniques, we obtain uniform L^p estimates for the Dirichlet, Neumann and regularity problems on bounded Lipschitz domains for a family of second order elliptic operators arising in the theory of homogenization. The results on the Neumann and regularity problems are new even for smooth domains.

MSC 2000: 35J25.

Keywords: Boundary value problem; Periodic coefficient; Lipschitz domain.

1 Introduction

Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ be a second order elliptic operator defined in $\mathbb{R}^{d+1} = \{X = (x, t) \in \mathbb{R}^d \times \mathbb{R}\}$, $d \geq 2$. We will always assume that the $(d+1) \times (d+1)$ coefficient matrix

$$A = A(X) = (a_{ij}(X)) \text{ is real and symmetric,} \quad (1.1)$$

and satisfies the ellipticity condition,

$$\mu|\xi|^2 \leq a_{ij}(X)\xi_i\xi_j \leq \frac{1}{\mu}|\xi|^2 \quad \text{for all } X, \xi \in \mathbb{R}^{d+1}, \quad (1.2)$$

*Supported in part by NSF grant DMS-0456583

†Supported in part by NSF grant DMS-0855294

where $\mu > 0$. In this paper we shall be interested in boundary value problems for $\mathcal{L}(u) = 0$ in the upper half-space $\mathbb{R}_+^{d+1} = \mathbb{R}^d \times (0, \infty)$ with L^p boundary data, under the assumption that the coefficients are periodic in the t variable,

$$A(x, t+1) = A(x, t) \quad \text{for } (x, t) \in \mathbb{R}^{d+1}. \quad (1.3)$$

More precisely, we will study the solvabilities of the L^p Dirichlet problem $(D)_p$

$$\begin{cases} \mathcal{L}(u) = 0 & \text{in } \Omega = \mathbb{R}_+^{d+1}, \\ u = f \in L^p(\partial\Omega) & \text{n.t. on } \partial\Omega \text{ and } (u)^* \in L^p(\partial\Omega), \end{cases} \quad (1.4)$$

the L^p Neumann problem $(N)_p$

$$\begin{cases} \mathcal{L}(u) = 0 & \text{in } \Omega = \mathbb{R}_+^{d+1}, \\ \frac{\partial u}{\partial \nu} = f \in L^p(\partial\Omega) & \text{on } \partial\Omega \text{ and } N(\nabla u) \in L^p(\partial\Omega), \end{cases} \quad (1.5)$$

where $\frac{\partial u}{\partial \nu}$ denote the conormal derivative associated with operator $\mathcal{L}$, and the L^p regularity problem $(R)_p$

$$\begin{cases} \mathcal{L}(u) = 0 & \text{in } \Omega = \mathbb{R}_+^{d+1}, \\ u = f \in \dot{W}^{1,p}(\partial\Omega) & \text{n.t. on } \partial\Omega \text{ and } N(\nabla u) \in L^p(\partial\Omega). \end{cases} \quad (1.6)$$

Here $(u)^*$ denotes the usual nontangential maximal function of u and $N(\nabla u)$ a generalized nontangential maximal function of ∇u . By $u = f$ n.t. on $\partial\Omega$ we mean that $u(X)$ converges to $f(P)$ as $X \rightarrow P$ nontangentially for a.e. $P \in \partial\Omega$. Under the periodic condition (1.3) as well as some (necessary) local solvability conditions on $\mathcal{L}$, we will show that the L^p Dirichlet problem is uniquely solvable for $2 - \delta < p < \infty$, and the L^p Neumann and regularity problems are uniquely solvable for $1 < p < 2 + \delta$. Furthermore, the solution to the Dirichlet problem satisfies the estimate $\|(u)^*\|_p \leq C\|u\|_p$, while the solutions to the L^p Neumann and regularity problems satisfy $\|N(\nabla u)\|_p \leq C\|\frac{\partial u}{\partial \nu}\|_p$ and $\|N(\nabla u)\|_p \leq C\|\nabla_x u\|_p$, respectively. These results extend the work on the L^p boundary value problems in $\mathbb{R}_+^{d+1}$ for second order elliptic operators with t -independent coefficients (or in bounded star-like Lipschitz domains for operators with radially independent coefficients). As we shall discuss later, since the local solvability conditions may be deduced from certain minimal smoothness conditions in the t -variable and our periodic condition is only imposed on the t -variable, the L^p global estimates on $(u)^*$ and $N(\nabla u)$ extend directly from $\mathbb{R}_+^{d+1}$ to the regions above Lipschitz graphs. As a consequence, by well known localization techniques, we obtain uniform estimates for the L^p boundary value problems in bounded Lipschitz domains for a family of second order elliptic operators arising in the theory of homogenization.

We should point out that the result mentioned above for the Dirichlet problem is in fact due to the late B. Dahlberg [9] (unpublished, personal communication). His proof, which is given in the Appendix for the sake of reference, depends on the ingenious use of Green's functions and harmonic measures. As in the case of his celebrated theorem on the Dirichlet problem for $\Delta u = 0$ in Lipschitz domains [10, 11], Dahlberg's method does not extend to the Neumann and regularity problems. Motivated by the work of Jerison and Kenig [16, 17, 18], we then seek to establish the global L^2 Rellich type estimate,

$$\left\| \frac{\partial u}{\partial \nu} \right\|_2 \approx \|\nabla_x u\|_2 \quad (1.7)$$

for suitable solutions of elliptic operators with t -periodic coefficients. In Section 3 we develop some new integral identities which play a crucial role in the proof of (1.7) and which may be regarded as the Rellich identities for operators with t -periodic coefficients. Indeed, in the case of constant or t -independent coefficients, Rellich identities are usually derived by using integration by parts on a form involving $\frac{\partial u}{\partial t}$. The basic insight here is to replace the t derivative of u by the difference $Q(u)(x, t) = u(x, t+1) - u(x, t)$. The t -periodicity of $A(x, t)$ is used in the fact that Q commutes with $\mathcal{L}$. In particular, $Q(u)$ is a solution whenever u is a solution. We further remark that these integral identities allow us to control the near-boundary integral

$$\int_0^1 \int_{\mathbb{R}^d} |\nabla u(x, t)|^2 dx dt, \quad (1.8)$$

by the integral of $|\frac{\partial u}{\partial \nu}|^2$ or $|\nabla_x u|^2$ on $\mathbb{R}^d$. The desired estimate (1.7) then follows by local solvability conditions.

With (1.7) at our disposal, we are able to give another proof of Dahlberg's theorem on the Dirichlet problem in Section 4 and more importantly, solve the L^2 regularity problem in Section 5 and the L^2 Neumann problem in Section 8. Furthermore, using the L^2 estimates and following the approach developed in [13] and [21], we establish the solvability of the L^p regularity and Neumann problems for $1 < p < 2$ in Sections 6 and 9. Although the range $2 < p < 2 + \delta$ may be also treated by the real variable arguments used in [13, 21], we choose to use a relatively new real variable approach, based on the weak reverse Hölder inequalities [25, 26]. In particular it allows us to show that for elliptic operators with coefficients satisfying (1.1)-(1.2), if the L^2 regularity problem $(R)_2$ is solvable and $\frac{1}{p} + \frac{1}{q} = 1$, then the Dirichlet problem $(D)_q$ and the regularity problem $(R)_p$ are equivalent.

There exists an extensive literature on boundary value problems with minimal smoothness assumptions on the coefficients or on the boundary of the domain in question. In particular the L^p Dirichlet, regularity and Neumann problems for $\Delta u = 0$ in Lipschitz domains, which are closely related to the problems we investigate here, were well understood thanks to [10, 11, 16, 17, 18, 29, 13]. Deep results on the L^p Dirichlet problem for a general second order elliptic equation $\mathcal{L}(u) = 0$ may be found in [8, 12, 14, 15] (see [19] for further references). The L^p regularity and Neumann problems for $\mathcal{L}(u) = 0$ were formulated and studied in [21, 22]. If the coefficient matrix A is t -independent, the L^p solvability of the Neumann and regularity problems in the upper half-space was essentially established in [21] for the sharp range $1 < p < 2 + \delta$, although the paper only treats the case of radially independent coefficients in the unit ball. We also mention that the L^2 boundary value problems for some operators with t -independent complex coefficients was recently studied by the method of layer potentials in [1]. Related work on L^p boundary value problems for operators with t -independent, but non-symmetric coefficients may be found in [20, 23]. Regarding the conditions on the t variable, it is worth mentioning that the global estimate $\|(u)^*\|_p \leq C\|f\|_p$ may fail for any p , even if the matrix $A(x, t)$ satisfies (1.1)-(1.2) and is C^∞ in $\mathbb{R}^{d+1}$.

Since the local solvability condition on $\mathcal{L}$ may be deduced from certain minimal smoothness assumption in the t variable on the coefficient matrix A , as we alluded to earlier, Dahlberg's theorem on the Dirichlet problem and our results on the Neumann and regularity problems extend readily to the case where

$$D = \{(x, t) \in \mathbb{R}^{d+1} : t > \psi(x)\} \quad (1.9)$$

is the region above a Lipschitz graph, using the bi-Lipschitzian map $(x, t) \rightarrow (x, t - \psi(x))$ from D to $\mathbb{R}_+^{d+1}$. Indeed, let

$$\eta(\rho) = \sup \{ |A(x, s_1) - A(x, s_2)| : x \in \mathbb{R}^d \text{ and } |s_1 - s_2| \leq \rho \} \quad (1.10)$$

and assume that

$$\int_0^1 \frac{(\eta(\rho))^2}{\rho} d\rho < \infty. \quad (1.11)$$

The following three theorems are proved in Section 10. The constant δ in Theorems 1.1, 1.2 and 1.3 depends only on $d, \mu, \eta(\rho)$ and $\|\nabla_x \psi\|_\infty$.

Theorem 1.1. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ with coefficient matrix satisfying (1.1), (1.2), (1.3) and (1.11). Let D be given by (1.9). Then there exists $\delta \in (0, 1)$ such that for any $f \in L^p(\partial D)$ with $2 - \delta < p < \infty$, there exists a unique solution to $\mathcal{L}(u) = 0$ in D with the property that $(u)^* \in L^p(\partial D)$ and $u = f$ n.t. on ∂D . Moreover, the solution u satisfies the estimate $\|(u)^*\|_p \leq C_p \|f\|_p$, where C_p depends only on $d, p, \mu, \eta(\rho)$ and $\|\nabla_x \psi\|_\infty$.*

Theorem 1.2. *Under the same assumptions on $\mathcal{L}$ and D as in Theorem 1.1, there exists $\delta > 0$ such that given any $g \in L^p(\partial D)$ with $1 < p < 2 + \delta$, there exists a solution, unique up to constants, to $\mathcal{L}(u) = 0$ in D such that $N(\nabla u) \in L^p(\partial D)$ and $\frac{\partial u}{\partial \nu} = g$ on ∂D . Moreover, the solution satisfies the estimate $\|N(\nabla u)\|_p \leq C_p \|g\|_p$, where C_p depends only on $d, p, \mu, \eta(\rho)$ and $\|\nabla_x \psi\|_\infty$.*

Let $\nabla_{\tan} u$ denote the tangential derivatives of u on $\partial\Omega$. We say $f \in \dot{W}^{1,p}(\partial\Omega)$ if $f \in L^p_{loc}(\partial\Omega)$ and $\nabla_{\tan} f \in L^p(\partial\Omega)$.

Theorem 1.3. *Under the same assumptions on $\mathcal{L}$ and D as in Theorem 1.1, there exists $\delta > 0$ such that given any $f \in \dot{W}^{1,p}(\partial D)$ with $1 < p < 2 + \delta$, there exists a unique solution to $\mathcal{L}(u) = 0$ in D such that $N(\nabla u) \in L^p(\partial D)$ and $u = f$ n.t. on ∂D . Moreover, the solution satisfies the estimate $\|N(\nabla u)\|_p \leq C_p \|\nabla_{\tan} f\|_p$, where C_p depends only on $d, p, \mu, \eta(\rho)$ and $\|\nabla_x \psi\|_\infty$.*

As we indicated earlier, Theorem 1.1 as well as Theorem 1.4 below are due to B. Dahlberg [9]. However, under the t -periodic condition, Theorems 1.2 and 1.3 are new even for operators with smooth coefficients on smooth domains. We further point out that by well known localization techniques, Theorems 1.1, 1.2 and 1.3 allow one to establish uniform estimates for the L^p boundary value problems in Lipschitz domains for a family of second order elliptic operators arising in the theory of homogenization (see e.g. [7]).

More precisely, we consider

$$\mathcal{L}_\varepsilon = -\operatorname{div} \left(A \left(\frac{X}{\varepsilon} \right) \nabla \right), \quad \varepsilon > 0, \quad (1.12)$$

where the matrix $A(X)$ is periodic with respect to the standard lattice,

$$A(X + Z) = A(X) \quad \text{for any } X \in \mathbb{R}^{d+1}, Z \in \mathbb{Z}^{d+1}. \quad (1.13)$$

In view of (1.11), we will assume that $A(X)$ is continuous and its modulus of continuity satisfies

$$\int_0^1 \frac{(\omega(\rho))^2}{\rho} d\rho < \infty, \quad (1.14)$$

where $\omega(\rho) = \sup \{|A(X) - A(Y)| : X, Y \in \mathbb{R}^{d+1} \text{ and } |X - Y| \leq \rho\}$.

The proofs of the following three theorems are given in Section 11.

Theorem 1.4. *Assume that the coefficient matrix $A(X)$ is real, symmetric, and satisfies the ellipticity condition (1.2), the periodicity condition (1.13) and the smoothness condition (1.14). Let Ω be a bounded Lipschitz domain in $\mathbb{R}^{d+1}$. Then there exist constants $\delta \in (0, 1)$ and $C > 0$ independent of ε , such that if $f \in L^p(\partial\Omega)$ for some $2 - \delta < p < \infty$, then the unique solution to the L^p Dirichlet problem: $\mathcal{L}_\varepsilon(u_\varepsilon) = 0$ in Ω , $u_\varepsilon = f$ n.t. on $\partial\Omega$, $(u_\varepsilon)^* \in L^p(\partial\Omega)$, satisfies the estimate $\|(u_\varepsilon)^*\|_p \leq C\|f\|_p$.*

Let $\frac{\partial u}{\partial \nu_\varepsilon}$ denote the conormal derivative associated with the operator $\mathcal{L}_\varepsilon$.

Theorem 1.5. *Let Ω be a bounded Lipschitz domain in $\mathbb{R}^{d+1}$. Assume that $A(X)$ satisfies the same assumption as in Theorem 1.4. Then there exist constants $\delta \in (0, 1)$ and $C > 0$ independent of ε , such that given any $g \in L^p(\partial\Omega)$ with $1 < p < 2 + \delta$ and mean value zero, the unique solution to the L^p Neumann problem: $\mathcal{L}_\varepsilon(u_\varepsilon) = 0$ in Ω , $\frac{\partial u_\varepsilon}{\partial \nu_\varepsilon} = g$ on $\partial\Omega$, $N(\nabla u_\varepsilon) \in L^p(\partial\Omega)$, satisfies the estimate $\|N(\nabla u_\varepsilon)\|_p \leq C\|g\|_p$.*

Theorem 1.6. *Let Ω be a bounded Lipschitz domain in $\mathbb{R}^{d+1}$. Assume that $A(X)$ satisfies the same assumption as in Theorem 1.4. Then there exist constants $\delta \in (0, 1)$ and $C > 0$ independent of $\varepsilon > 0$, such that given any $f \in W^{1,p}(\partial\Omega)$ with $1 < p < 2 + \delta$, the unique solution to the L^p regularity problem: $\mathcal{L}_\varepsilon(u_\varepsilon) = 0$ in Ω , $u_\varepsilon = f$ n.t. on $\partial\Omega$, $N(\nabla u_\varepsilon) \in L^p(\partial\Omega)$ satisfies the estimate $\|N(\nabla u_\varepsilon)\|_p \leq C \{ \|\nabla_{\tan} f\|_p + |\partial\Omega|^{\frac{1}{1-d}} \|f\|_p \}$.*

We remark that Dahlberg's work on operators with periodic coefficients was inspired by a series of remarkable papers [2, 3, 4, 5, 6] of M. Avellaneda and F. Lin on elliptic homogenization problems. In particular, the uniform estimate $\|(u_\varepsilon)^*\|_p \leq C\|f\|_p$ in Theorem 1.4 for $1 < p < \infty$ was established on $C^{1,\gamma}$ domains by Avellaneda and Lin in [2, 3, 6] for second order elliptic equations and systems in divergence form with C^α periodic coefficients. Related work on the boundedness of Riesz transforms $\nabla(\mathcal{L}_\varepsilon)^{-1/2}$ on Lipschitz and C^1 domains may be found in [28]. However, to the authors' best knowledge, the uniform L^p estimates for periodic operators have not been studied before for the regularity and Neumann problems; Theorems 1.5 and 1.6 are new even for smooth domains.

Finally we mention that as in the case of operators with constant coefficients, the Rellich estimates we develop in this paper, can be used to solve the L^p boundary value problems by the method of layer potentials. This would enable us to represent the solutions in Theorems 1.4, 1.5 and 1.6 in terms of layer potentials with density functions uniformly bounded in appropriate spaces. Even more importantly, the method of layer potentials can be applied to elliptic systems and would allow us to establish uniform L^p estimates for elliptic systems with periodic coefficients in nonsmooth domains. This line of ideas will be fully developed in a forthcoming paper [24].

Acknowledgment. The first named author is indebted to the late Björn Dahlberg for sharing with him the proof of Theorem 4.1 on the Dirichlet problem. Our work was motivated in part by Dahlberg's theorem. Dahlberg died suddenly on January 30, 1998 and did not publish his proof. As we mentioned earlier, we include a version of his original proof in the Appendix for future reference. We dedicate this paper to the memory of Björn Dahlberg.

2 Notations and Preliminaries

We will use X, Y, Z to denote points in $\mathbb{R}^{d+1}$ and x, y, z points in $\mathbb{R}^d$. Balls in $\mathbb{R}^d$ are denoted by $B(x, r) = \{y \in \mathbb{R}^d : |y - x| < r\}$. By $T(x, r)$ we mean the cylinder $B(x, r) \times (0, r)$. We will use ∇ for the gradient in $\mathbb{R}^{d+1}$ and ∇_x for the gradient in $\mathbb{R}^d$.

The classical Dirichlet problem in $\Omega = \mathbb{R}_+^{d+1}$.

Let $f \in C(\mathbb{R}^d)$ such that $f(x) \rightarrow 0$ as $|x| \rightarrow \infty$. Then there exists a unique weak solution to $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ such that $u \in C(\mathbb{R}_+^{d+1} \times [0, \infty))$, $u(x, 0) = f(x)$ for $x \in \mathbb{R}^d$ and u is bounded in $\mathbb{R}_+^{d+1}$. Moreover, the solution satisfies $\sup\{|u(x, t)| : x \in \mathbb{R}^d, t > 0\} \leq \|f\|_\infty$. The uniqueness follows from Liouville's Theorem by a reflection argument. To establish the existence, one may assume that $f \geq 0$. Let $\Omega_k = T(0, k) = B(0, k) \times (0, k)$ and φ a continuous decreasing function on $[0, \infty)$ such that $0 \leq \varphi \leq 1$, $\varphi(s) = 1$ for $0 \leq s \leq (1/2)$ and $\varphi(s) = 0$ for $s > (3/4)$. Let u_k be the solution to the classical Dirichlet problem for $\mathcal{L}(u) = 0$ in $T(0, k)$ with boundary data $f(x)\varphi(|x|/k)$ on $\{(x, 0) : |x| < k\}$ and zero otherwise. Since $u_k \leq u_{k+1}$ on Ω_k and $\|u_k\|_{L^\infty(\Omega_k)} \leq \|f\|_\infty$, it follows that $u(x) = \lim_{k \rightarrow \infty} u_k(x)$ exists and is finite for any $x \in \mathbb{R}_+^{d+1}$ and $\|u\|_{L^\infty(\mathbb{R}_+^{d+1})} \leq \|f\|_\infty$. Also, we may deduce from the estimate

$$\sup_{\Omega_\ell} |u_k - u_m| \leq C(u_k - u_m)(0, \ell/2), \quad \text{for } k > m > \ell$$

that as $k \rightarrow \infty$, u_k converges to u uniformly on $\overline{\Omega_\ell}$ for any $\ell > 1$. This implies that $\mathcal{L}(u) = 0$ in $\Omega = \mathbb{R}_+^{d+1}$, u is continuous in $\overline{\Omega}$ and $u(x, 0) = f(x)$.

By the Riesz representation theorem, there exists a family of regular Borel measures $\{\omega^X : X \in \mathbb{R}_+^{d+1}\}$ on $\mathbb{R}^d$, called $\mathcal{L}$ -harmonic measures, such that

$$u(X) = \int_{\mathbb{R}^d} f(y) d\omega^X(y). \quad (2.1)$$

Let v_k be the solution to the classical Dirichlet problem in $\mathbb{R}_+^{d+1}$ with data $\varphi(|x|/k)$. It follows from the uniqueness that $v_k(X) \rightarrow 1$ as $k \rightarrow \infty$ for any $X \in \mathbb{R}_+^{d+1}$. This shows that ω^X is a probability measure, as in the case of bounded domains. Furthermore, let ω_k^X denote the $\mathcal{L}$ -harmonic measure for the cylinder $B(0, k) \times (0, k)$. It is not hard to see that $\omega_k^X(B) \rightarrow \omega^X(B)$ as $k \rightarrow \infty$, for any ball $B \subset \mathbb{R}^d$ and $X \in \mathbb{R}_+^{d+1}$.

Green's function on $\Omega = \mathbb{R}_+^{d+1}$.

Let $G_k(X, Y)$ denote the Green's function for $\mathcal{L}$ on the domain $\Omega_k = T(0, k)$. Since $0 \leq G_k(X, Y) \leq G_{k+1}(X, Y)$ for $X, Y \in \Omega_k$, the Green's function for $\mathcal{L}$ on $\Omega = \mathbb{R}_+^{d+1}$ may be defined by

$$G(X, Y) = \lim_{k \rightarrow \infty} G_k(X, Y) \quad \text{for } X, Y \in \mathbb{R}_+^{d+1} \text{ and } X \neq Y.$$

By well known properties of G_k , we have

$$0 \leq G(X, Y) \leq \frac{C}{|X - Y|^{d-1}} \quad \text{and} \quad G(X, Y) \leq \frac{Ct^\alpha}{|X - Y|^{d-1+\alpha}} \quad (2.2)$$

for $X = (x, t)$, $Y \in \mathbb{R}_+^{d+1}$, where C and α are positive constants depending only on d and μ . Since $\omega_k^X(B(y, r)) \approx r^{d-1}G_k(X, (y, r))$ for any $X \in \Omega_k \setminus T(y, 2r)$ [8], by letting $k \rightarrow \infty$, we obtain

$$\omega^X(B(y, r)) \approx r^{d-1}G(X, (y, r)) \quad \text{for any } X \notin T(y, 2r). \quad (2.3)$$

Two properties of nonnegative weak solutions.

Let u be a nonnegative weak solution of $\mathcal{L}(u) = 0$ in $T(x_0, 2r)$ for some $x_0 \in \mathbb{R}^d$ and $r > 0$. Suppose that $u \in C(\overline{T(x_0, 2r)})$ and $u(x, 0) = 0$ on $B(x_0, 2r)$. Then we have the boundary Harnack inequality,

$$u(x, t) \leq Cu(x_0, r) \quad \text{for any } (x, t) \in T(x_0, r). \quad (2.4)$$

We will also need the following comparison principle,

$$\frac{u(x, t)}{v(x, t)} \leq C \frac{u(x_0, r)}{v(x_0, r)} \quad \text{for any } (x, t) \in T(x_0, r), \quad (2.5)$$

where u, v are two nonnegative weak solutions in $T(x_0, 2r)$ such that $u, v \in C(\overline{T(x_0, 2r)})$ and $u(x, 0) = v(x, 0) = 0$ on $B(x_0, 2r)$ (see e.g. [8]). The constants in (2.4) and (2.5) depend only on d and μ .

Difference operator Q .

Define

$$Q(u)(x, t) = u(x, t+1) - u(x, t) \quad \text{for } (x, t) \in \mathbb{R}^{d+1}. \quad (2.6)$$

Clearly, $Q(\frac{\partial u}{\partial x_i}) = \frac{\partial}{\partial x_i}Q(u)$ for $i = 1, \dots, d+1$. It is also easy to verify that

$$Q(uv) - Q(u)Q(v) = uQ(v) + vQ(u) \quad (2.7)$$

and

$$\iint_{T(x, t)} Q(u)(y, s) dy ds = \int_t^{t+1} \int_{B(x, t)} u(y, s) dy ds - \int_0^1 \int_{B(x, t)} u(y, s) dy ds. \quad (2.8)$$

Nontangential maximal functions.

For a function u defined in $\mathbb{R}_+^{d+1}$, we let

$$N(u)(x) = \sup \left\{ \left(\frac{1}{t|B(x, t)|} \int_{\frac{t}{2}}^{\frac{3t}{2}} \int_{B(x, t)} |u(y, s)|^2 dy ds \right)^{1/2} : 0 < t < \infty \right\}. \quad (2.9)$$

This is a variant of the usual nontangential maximal function $(u)^*$ which is defined by

$$(u)^*(x) = \sup \{ |u(y, t)| : 0 < t < \infty \text{ and } |y - x| < t \}. \quad (2.10)$$

Clearly, $N(u)(x) \leq (u)^*(x)$. If u is a weak solution to $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ or any function that has the property

$$|u(x, t)| \leq C \left(\frac{1}{r^{d+1}} \iint_{B((x, t), r)} |u(y, s)|^2 dy ds \right)^{1/2},$$

whenever $B((x, t), r) \subset \mathbb{R}_+^{d+1}$, then $\|N(u)\|_p \approx \|(u)^*\|_p$ for any $0 < p < \infty$. We also need to introduce $N_r(u)$ and $(u)_r^*$ which are defined by restricting the variable t in (2.9) and (2.10) respectively to $0 < t < r/2$. The definitions of $N(u)$ and $(u)^*$ extend naturally to regions above Lipschitz graphs and to bounded Lipschitz domains.

We conclude this section with a lemma which allows us to estimate $\|N(\nabla u)\|_p$ by $\|N_4(\nabla u)\|_p$ and $\|(Q(u))^*\|_p$.

Lemma 2.1. *Let $u \in C^1(\overline{T(0, 3R)})$ be a solution to $\mathcal{L}(u) = 0$ in $T(0, 3R)$ for some $R > 10$. Let $f(x) = u(x, 0)$. Then for any $x \in B(0, R)$,*

$$N_R(\nabla u)(x) \leq C M \left([N_4(\nabla u) + |\nabla_x f| + (Q(u))_R^*] \chi_{B(0, 3R)} \right)(x), \quad (2.11)$$

where M denotes the Hardy-Littlewood maximal operator on $\mathbb{R}^d$ and C depends only on d and μ .

Proof. We begin by fixing $(x_0, t_0) \in \mathbb{R}^{d+1}$ with $x_0 \in B(0, R)$ and $t_0 \in (2, R/2)$. Note that by Cacciopoli's inequality and interior regularity of weak solutions,

$$\begin{aligned} & \left(\frac{1}{t_0^{d+1}} \int_{|s-t_0| < \frac{t_0}{2}} \int_{B(x_0, t_0)} |\nabla u(y, s)|^2 dy ds \right)^{1/2} \\ & \leq \frac{C}{t_0} \left(\frac{1}{t_0^{d+1}} \int_{|s-t_0| < \alpha_1 t_0} \int_{B(x_0, \beta_1 t_0)} |u(y, s) - E|^2 dy ds \right)^{1/2} \\ & \leq \frac{C}{t_0^{d+2}} \int_{|s-t_0| < \alpha_2 t_0} \int_{B(x_0, \beta_2 t_0)} |u(y, s) - E| dy ds, \end{aligned} \quad (2.12)$$

where $(1/2) < \alpha_1 < \alpha_2 < (3/4)$, $1 < \beta_1 < \beta_2 < (3/2)$ and $E \in \mathbb{R}$. To estimate the right-hand side of (2.12), we use

$$|u(y, s) - E| \leq |u(y, s) - u(y, 0)| + |u(y, 0) - E|$$

and choose E to be the average of $u(y, 0)$ over $B(x_0, \beta_1 t_0)$. By Poincaré's inequality, we have

$$\begin{aligned} & \frac{1}{t_0^{d+2}} \int_{|s-t_0| < \alpha_2 t_0} \int_{B(x_0, \beta_2 t_0)} |u(y, 0) - E| dy ds \\ & \leq \frac{C}{t_0^d} \int_{B(x_0, \beta_2 t_0)} |\nabla_x f| dx \leq CM(|\nabla_x f| \chi_{B(0, 3R)})(x_0). \end{aligned} \quad (2.13)$$

Next we observe that for any $s \in (0, 2t_0)$,

$$|u(y, s) - u(y, 0)| \leq Ct_0(Q(u))_R^*(y) + \int_0^1 |\nabla u(y, t)| dt. \quad (2.14)$$

In view of (2.12), the first term in the right-hand side of (2.14) can be handled easily by $M((Q(u))_R^* \chi_{B(0,3R)})(x_0)$. Thus it remains to estimate

$$\frac{1}{t_0^{d+1}} \int_{B(x_0, \beta_2 t_0)} \int_0^1 |\nabla u(y, t)| dt dy.$$

This may be done by observing that

$$C \int_{B(x_0, 4t_0)} N_4(\nabla u) dx \geq \int_0^1 \int_{B(x_0, 2t_0)} |\nabla u(y, s)| dy ds.$$

□

Remark 2.2. It follows from (2.11) that for $1 < p < \infty$,

$$\begin{aligned} \int_{B(0, R)} |N_R(\nabla u)|^p dx &\leq C \int_{B(0, 3R)} |\nabla_x u(x, 0)|^p dx + C \int_{B(0, 3R)} |N_4(\nabla u)|^p dx \\ &\quad + C \int_{B(0, 3R)} |(Q(u))_R^*|^p dx \end{aligned} \quad (2.15)$$

for any $R > 10$. Thus, if $u \in C^1(\mathbb{R}^d \times [0, \infty))$ and $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$, we may let $R \rightarrow \infty$ in (2.15) to obtain

$$\begin{aligned} \int_{\mathbb{R}^d} |N(\nabla u)|^p dx &\leq C \int_{\mathbb{R}^d} |\nabla_x u(x, 0)|^p dx + C \int_{\mathbb{R}^d} |N_4(\nabla u)|^p dx \\ &\quad + C \int_{\mathbb{R}^d} |(Q(u))^*|^p dx, \end{aligned} \quad (2.16)$$

where C depends only on d, p and μ .

3 Integral identities

In this section we develop some new integral identities that will play a key role in our approach to the L^2 boundary value problems in $\mathbb{R}_+^{d+1}$ for elliptic equations with t -periodic coefficients.

Lemma 3.1. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ with coefficients satisfying conditions (1.1)-(1.2). Let $\Omega \subset \mathbb{R}^{d+1}$ be a bounded Lipschitz domain and $u \in W^{1,2}(\Omega)$ a variational solution to the Neumann problem: $\mathcal{L}(u) = 0$ in Ω , $\frac{\partial u}{\partial \nu} = g \in L^2(\partial\Omega)$. Assume that $u \in W^{1,2}(\Omega_1)$ where Ω_1 contains Ω and $\{(x, t+1) : (x, t) \in \Omega\}$. Then*

$$\begin{aligned} & - \int_{\Omega} a_{ij} Q \left(\frac{\partial u}{\partial x_i} \cdot \frac{\partial u}{\partial x_j} \right) dX + \int_{\Omega} a_{ij} Q \left(\frac{\partial u}{\partial x_i} \right) \cdot Q \left(\frac{\partial u}{\partial x_j} \right) dX \\ & = -2 \int_{\partial\Omega} g Q(u) d\sigma, \end{aligned} \quad (3.1)$$

where indices i, j are summed from 1 to $d+1$ and we identify x_{d+1} with t .

Proof. Since $Q(u) \in W^{1,2}(\Omega)$, it follows from the definition of variational solutions that

$$\begin{aligned} \int_{\partial\Omega} g Q(u) d\sigma &= \int_{\Omega} a_{ij} \frac{\partial u}{\partial x_j} \cdot Q\left(\frac{\partial u}{\partial x_i}\right) dX \\ &= \frac{1}{2} \int_{\Omega} a_{ij} \left\{ \frac{\partial u}{\partial x_j} \cdot Q\left(\frac{\partial u}{\partial x_i}\right) + \frac{\partial u}{\partial x_i} \cdot Q\left(\frac{\partial u}{\partial x_j}\right) \right\} dX, \end{aligned}$$

where we have used the symmetry condition $a_{ij} = a_{ji}$ in the second equation. In view of (2.7), this gives

$$\int_{\partial\Omega} g Q(u) d\sigma = \frac{1}{2} \int_{\Omega} a_{ij} \left\{ Q\left(\frac{\partial u}{\partial x_i}\right) \cdot \frac{\partial u}{\partial x_j} - Q\left(\frac{\partial u}{\partial x_j}\right) \cdot \frac{\partial u}{\partial x_i} \right\} dX,$$

from which the desired identity follows. $\square$

In the rest of this section we will assume that a_{ij} satisfy the periodic condition (1.3). To justify the use of integration by parts, we will also assume that both a_{ij} and u are sufficiently smooth. However all constants C in this section depend only on d and μ .

The following lemma is crucial in our argument for the L^2 Neumann problem.

Lemma 3.2. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ with C^∞ coefficients satisfying (1.1), (1.2) and (1.3). Let $R > 10$ and $u \in C^1(\overline{\Omega})$ be a solution to $\mathcal{L}(u) = 0$ in $\Omega = T(x_0, 3R)$ and $\frac{\partial u}{\partial \nu} = g$ on $B(x_0, 3R) \times \{0\}$. Then*

$$\int_0^1 \int_{B(x_0, R)} |\nabla u|^2 dx dt \leq C \int_{B(x_0, 2R)} |g|^2 dx + \frac{C}{R} \iint_{T(x_0, 3R) \setminus T(x_0, R)} |\nabla u|^2 dx dt, \quad (3.2)$$

where C depends only on d and μ .

Proof. We may assume that $x_0 = 0$. Let $\Omega_r = T(0, r) = B(0, r) \times (0, r)$ for $r \in (R, 2R)$. By the periodicity assumption (1.3) and (2.8), we have

$$\begin{aligned} & - \iint_{\Omega_r} a_{ij} Q\left(\frac{\partial u}{\partial x_i}\right) \cdot \frac{\partial u}{\partial x_j} dx dt = - \iint_{\Omega_r} Q\left(a_{ij} \frac{\partial u}{\partial x_i}\right) \cdot \frac{\partial u}{\partial x_j} dx dt \\ & = \int_0^1 \int_{B(0, r)} a_{ij} \frac{\partial u}{\partial x_i} \cdot \frac{\partial u}{\partial x_j} dx dt - \int_r^{r+1} \int_{B(0, r)} a_{ij} \frac{\partial u}{\partial x_i} \cdot \frac{\partial u}{\partial x_j} dx dt. \end{aligned}$$

It then follows from (3.1) that

$$\begin{aligned} & \int_0^1 \int_{B(0, r)} a_{ij} \frac{\partial u}{\partial x_i} \cdot \frac{\partial u}{\partial x_j} dx dt + \iint_{\Omega_r} a_{ij} Q\left(\frac{\partial u}{\partial x_i}\right) Q\left(\frac{\partial u}{\partial x_j}\right) dx dt \\ & = -2 \int_{\partial\Omega_r} \frac{\partial u}{\partial \nu} \cdot Q(u) d\sigma + \int_r^{r+1} \int_{B(0, r)} a_{ij} \frac{\partial u}{\partial x_i} \cdot \frac{\partial u}{\partial x_j} dx dt, \end{aligned} \quad (3.3)$$

for $r \in (R, 2R)$. By the Cauchy inequality, this leads to

$$\begin{aligned}
\int_0^1 \int_{B(0,r)} |\nabla u|^2 dx dt &\leq \delta \int_{B(0,r)} |Q(u)(x, 0)|^2 dx + C_\delta \int_{B(0,r)} |g(x)|^2 dx \\
&\quad + C \int_{B(0,r)} |\nabla u(x, r)|^2 dx + C \int_{B(0,r)} |Q(u)(x, r)|^2 dx \\
&\quad + C \int_r^{r+1} \int_{B(0,r)} |\nabla u|^2 dx dt + C \int_0^r \int_{\partial B(0,r)} |\nabla u|^2 d\sigma dt \\
&\quad + C \int_0^r \int_{\partial B(0,r)} |Q(u)|^2 d\sigma dt,
\end{aligned} \tag{3.4}$$

for any $0 < \delta < 1$. Using

$$\begin{aligned}
\int_{B(0,r)} |Q(u)(x, t)|^2 dx &\leq \int_t^{t+1} \int_{B(0,r)} |\nabla u|^2 dx ds, \\
\int_0^r \int_{\partial B(0,r)} |Q(u)|^2 d\sigma dt &\leq \int_0^{r+1} \int_{\partial B(0,r)} |\nabla u|^2 d\sigma dt,
\end{aligned}$$

we obtain from (3.4) that

$$\begin{aligned}
\int_0^1 \int_{B(0,R)} |\nabla u|^2 dx dt &\leq C \int_{B(0,2R)} |g(x)|^2 dx + C \int_{B(0,2R)} |\nabla u(x, r)|^2 dx \\
&\quad + C \int_r^{r+1} \int_{B(0,2R)} |\nabla u|^2 dx dt \\
&\quad + C \int_0^{3R} \int_{\partial B(0,r)} |\nabla u|^2 d\sigma dt,
\end{aligned} \tag{3.5}$$

for $r \in (R, 2R)$. The desired estimate now follows by integrating both sides of (3.5) with respect to r over the interval $(R, 2R)$. $\square$

Remark 3.3. Under the same assumption as in Lemma 3.2, we also have

$$\int_0^6 \int_{B(x_0,R)} |\nabla u|^2 dx dt \leq C \int_{B(x_0,2R)} |g|^2 dx + \frac{C}{R} \iint_{T(x_0,3R) \setminus T(x_0,R)} |\nabla u|^2 dx dt. \tag{3.6}$$

To see (3.6), one simply replaces $Q(u)$ with $\tilde{Q}(u) = u(x, t+6) - u(x, t)$ in the proof.

The following lemma is the key in our approach to the L^2 regularity problem.

Lemma 3.4. *Under the same assumptions on $\mathcal{L}$ as in Lemma 3.2, there exists $C = C(d, \mu) > 0$ such that*

$$\int_0^1 \int_{B(x_0,R)} |\nabla u|^2 dx dt \leq C \int_{B(x_0,2R)} |\nabla_x f|^2 dx + \frac{C}{R} \iint_{T(x_0,3R) \setminus T(x_0,R)} |\nabla u|^2 dx dt, \tag{3.7}$$

where $u \in C^1(\overline{T(x_0, 3R)}) \cap C^2(T(x_0, 3R))$ is a solution to $\mathcal{L}(u) = 0$ in $T(x_0, 3R)$ and $u = f$ on $B(x_0, 3R) \times \{0\}$, $R > 10$.

Proof. We may assume that $x_0 = 0$. As in the proof of Lemma 3.2, we let $\Omega_r = B(0, r) \times (0, r)$ for $r \in (R, 2R)$. By the periodicity assumption, $Q(u)$ is a solution in Ω_r . It follows that for $r \in (R, 2R)$,

$$\begin{aligned}
& \int_{\partial\Omega_r} \frac{\partial u}{\partial \nu} \cdot Q(u) d\sigma \\
&= \int_{\partial\Omega_r} u \cdot \frac{\partial}{\partial \nu} Q(u) d\sigma \\
&= \int_{\partial\Omega_r} u n_i \cdot \int_0^1 \frac{\partial}{\partial s} \left\{ a_{ij}(x, t+s) \frac{\partial u}{\partial x_j}(x, t+s) \right\} ds d\sigma(x, t) \\
&= \int_{\partial\Omega_r} u n_i \cdot \left\{ \frac{\partial}{\partial t} \int_0^1 a_{ij}(x, t+s) \frac{\partial u}{\partial x_j}(x, t+s) ds \right\} d\sigma(x, t) \\
&= \int_{\partial\Omega_r} u \left(n_i \frac{\partial}{\partial t} - n_{d+1} \frac{\partial}{\partial x_i} \right) \left\{ \int_0^1 a_{ij}(x, t+s) \frac{\partial u}{\partial x_j}(x, t+s) ds \right\} d\sigma(x, t) \\
&= - \int_{\partial\Omega_r} \left(n_i \frac{\partial}{\partial t} - n_{d+1} \frac{\partial}{\partial x_i} \right) u \cdot \left\{ \int_0^1 a_{ij}(x, t+s) \frac{\partial u}{\partial x_j}(x, t+s) ds \right\} d\sigma(x, t).
\end{aligned} \tag{3.8}$$

Consequently, by the Cauchy inequality, we obtain

$$\begin{aligned}
& \left| \int_{\partial\Omega_r} \frac{\partial u}{\partial \nu} \cdot Q(u) d\sigma \right| \\
&\leq C \int_0^1 \int_{\partial\Omega_r} |\nabla_{tan} u(x, t)| |\nabla u(x, t+s)| d\sigma(x, t) ds \\
&\leq \delta \int_0^1 \int_{B(0, r)} |\nabla u(x, t)|^2 dx dt + C_\delta \int_{B(0, r)} |\nabla_x u(x, 0)|^2 dx \\
&\quad + C \int_{B(0, r)} |\nabla u(x, r)|^2 dx + C \int_r^{r+1} \int_{B(0, r)} |\nabla u(x, t)|^2 dx dt \\
&\quad + C \int_0^{r+1} \int_{\partial B(0, r)} |\nabla u(x, t)|^2 d\sigma(x) dt,
\end{aligned} \tag{3.9}$$

for any $0 < \delta < 1$, where ∇_{tan} denotes the tangential gradient on $\partial\Omega_r$. This, together with (3.3), implies that

$$\begin{aligned}
& \int_0^1 \int_{B(0, R)} |\nabla u|^2 dx dt \\
&\leq C \int_{B(0, 2R)} |\nabla_x f|^2 dx + C \int_{B(0, 2R)} |\nabla u(x, r)|^2 dx \\
&\quad + C \int_0^{3R} \int_{\partial B(0, r)} |\nabla u|^2 d\sigma dt + C \int_r^{r+1} \int_{B(0, 2R)} |\nabla u|^2 dx dt,
\end{aligned} \tag{3.10}$$

for $r \in (R, 2R)$. The estimate (3.7) follows by integrating both sides of (3.10) with respect to r over the interval $(R, 2R)$. $\square$

Remark 3.5. By replacing $Q(u)$ with $\tilde{Q}(u)(x, t) = u(x, t+6) - u(x, t)$ in the proof of Lemma 3.4, we may obtain

$$\int_0^6 \int_{B(x_0, R)} |\nabla u|^2 dx dt \leq C \int_{B(x_0, 2R)} |\nabla_x f|^2 dx + \frac{C}{R} \iint_{T(x_0, 3R) \setminus T(x_0, R)} |\nabla u|^2 dx dt. \quad (3.11)$$

4 The L^p Dirichlet problem: a theorem of Dahlberg

To solve the L^p Dirichlet problem, we will impose a local solvability condition on $\mathcal{L}$: for any $x_0 \in \mathbb{R}^d$ and $0 < r \leq 1$,

$$\int_{B(x_0, r)} \limsup_{t \rightarrow 0} \left(\frac{u(x, t)}{t} \right)^2 dx \leq \frac{C_0}{r^3} \iint_{T(x_0, 2r)} |u(x, t)|^2 dx dt, \quad (4.1)$$

whenever $u \in W^{1,2}(T(x_0, 4r))$ is a nonnegative weak solution of $\mathcal{L}(u) = 0$ in $T(x_0, 4r)$ such that $u \in C(\overline{T(x_0, 4r)})$ and $u(x, 0) = 0$ on $B(x_0, 4r)$. Recall that for $Z \in \mathbb{R}_+^{d+1} \setminus T(x, 2t)$,

$$\frac{\omega^Z(B(x, t))}{|B(x, t)|} \approx \frac{G((x, t), Z)}{t}, \quad (4.2)$$

where ω^Z denotes the $\mathcal{L}$ -harmonic measure on $\mathbb{R}_+^{d+1}$ and $G(X, Z)$ the Green's function on $\mathbb{R}_+^{d+1}$. By applying the condition (4.1) to $u(X) = G(X, Z)$ for $Z \in \mathbb{R}_+^{d+1} \setminus T(x_0, 4r)$, we may deduce that ω^Z is absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}^d$. Furthermore, the kernel function

$$K(x, Z) := \lim_{t \rightarrow 0} \frac{\omega^Z(B(x, t))}{|B(x, t)|}$$

satisfies the reverse Hölder inequality

$$\left(\frac{1}{|B|} \int_B |K(y, Z)|^2 dy \right)^{1/2} \leq \frac{C}{|B|} \int_B K(y, Z) dy, \quad (4.3)$$

for $B = B(x, r)$ with $0 < r \leq 1$, if $Z \notin T(x, 2r)$. In fact, by the comparison principle (2.5), conditions (4.1) and (4.3) are equivalent.

The goal of this section is to show that with the additional periodic condition (1.3), the reverse Hölder inequality (4.3) holds without the restriction $r \leq 1$. As a consequence, the L^p Dirichlet problem for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable for $2 - \delta < p < \infty$, under the assumptions (1.1), (1.2), (1.3) and (4.1). The following theorem is due to B. Dahlberg [9].

Theorem 4.1. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ be an elliptic operator satisfying conditions (1.1), (1.2), (1.3) and (4.1). Then there exists $\delta = \delta(d, \mu, C_0) \in (0, 1)$ such that given any $f \in L^p(\mathbb{R}^d)$ with $2 - \delta < p < \infty$, there exists a unique weak solution to $\mathcal{L}(u) = 0$ in $\mathbb{R}^d$ with the property that $u = f$ n.t. on $\mathbb{R}^d$ and $(u)^* \in L^p(\mathbb{R}^d)$. Moreover the solution satisfies the estimate $\|(u)^*\|_p \leq C_p \|f\|_p$, where C_p depends only on d, p, μ and the constant C_0 in (4.1).*

The key step in the proof of Theorem 4.1 is the following lemma.

Lemma 4.2. *Under the same assumption on $\mathcal{L}$ as in Theorem 4.1, estimate (4.1) and hence (4.3) hold for $r > 1$.*

Proof. We will prove this lemma under the additional assumption that the coefficient matrix $A(X)$ is sufficiently smooth. The assumption allows us to use estimate (3.7). For the sake of completeness and future reference we shall provide a version of Dahlberg's original proof of Lemma 4.2 in the Appendix. We point out that our proofs of the theorems stated in the Introduction, given in Sections 10 and 11, involve an approximation argument and do not rely on Dahlberg's proof.

The case $1 < r \leq 10$ follows easily by covering $B(x_0, r)$ with balls of radius 1. If $r > 10$, we also cover $B(x_0, r)$ with a sequence of balls $\{B(x_k, 1)\}$ of radius 1 such that

$$B(x_0, r) \subset \bigcup_k B(x_k, 1) \subset B(x_0, r+1) \quad \text{and} \quad \sum_k \chi_{B(x_k, 1)} \leq C.$$

Let I denote the left-hand side of (4.1). It follows from the assumption (4.1) (with $r = 1$) and boundary Harnack inequality (2.4) that

$$\begin{aligned} I &\leq C \sum_k \int_{T(x_k, 2)} |u(X)|^2 dX \leq C \sum_k \int_{T(x_k, 1)} |u(X)|^2 dX \\ &\leq \sum_k \int_{T(x_k, 1)} |\nabla u(X)|^2 dX \leq C \int_0^1 \int_{B(x_0, r+1)} |\nabla u(x, t)|^2 dx dt. \end{aligned} \tag{4.4}$$

In view of Lemma 3.4, we obtain

$$\begin{aligned} I &\leq \frac{C}{r} \iint_{T(x_0, 3.5r)} |\nabla u|^2 dx dt \leq \frac{C}{r^3} \iint_{T(x_0, 3.6r)} |u|^2 dx dt \\ &\leq \frac{C}{r^3} \iint_{T(x, 2r)} |u|^2 dx dt, \end{aligned}$$

where we also used Cacciopoli's inequality as well as the Harnack inequality (2.4). $\square$

For $1 < p < \infty$, we say the L^p Dirichlet problem $(D)_p$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable if for any $f \in C_c(\mathbb{R}^d)$, the solution to the classical Dirichlet problem for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ with boundary data f satisfies the estimate $\|(u)^*\|_p \leq C\|f\|_p$.

Lemma 4.3. *Under the same assumption on $\mathcal{L}$ as in Theorem 4.1, the L^p Dirichlet problem for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable for $2 - \delta < p < \infty$. Moreover, we have $\|(u)^*\|_p \leq C\|f\|_p$ for $2 - \delta < p < \infty$, where $\delta = \delta(d, \mu, C_0) \in (0, 1)$ and $C = C(d, \mu, C_0, p) > 0$.*

Proof. By the self-improving property of the reverse Hölder inequality, there exists $q_0 > 2$ such that for $B = B(x_0, r)$ and $Z = (z, z_{d+1}) \in \mathbb{R}_+^{d+1} \setminus T(x_0, 4r)$,

$$\left\{ \frac{1}{|B|} \int_B |K(x, Z)|^{q_0} dx \right\}^{1/q_0} \leq \frac{C}{|B|} \int_B K(x, Z) dx = C \frac{\omega^Z(B)}{|B|} \leq \frac{C(z_{d+1})^\alpha}{r^{d+\alpha}}, \tag{4.5}$$

where we used (2.3) and (2.2). Since

$$u(Z) = \int_{\mathbb{R}^d} K(x, Z) f(x) dx,$$

one may deduce from (4.5) that

$$(u)^*(x) \leq C \{M(|f|^{p_0})(x)\}^{1/p_0}$$

for any $x \in \mathbb{R}^d$, where $p_0 = q'_0 < 2$ and M denotes the Hardy-Littlewood maximal operator on $\mathbb{R}^d$. This gives $\|(u)^*\|_p \leq C\|f\|_p$ for any $p > p_0 = 2 - \delta$. $\square$

We need the following Cacciopoli-type lemma in the proof of uniqueness.

Lemma 4.4. *Let u, v be two weak solutions in $B(x_0, 2R) \times (r/2, 3r)$ where $R \geq r$. Suppose that either u or v is nonnegative. Then*

$$\int_r^{2r} \int_{B(x_0, R)} |\nabla u| |v| \, dx dt \leq \frac{C}{r} \int_{\frac{r}{2}}^{3r} \int_{B(x_0, 2R)} |u| |v| \, dx dt,$$

where C depends only on d and μ .

Proof. Assume that $u \geq 0$. Let $X \in B(x_0, R) \times (r, 2r)$. Using Cauchy inequality and Cacciopoli's inequality, we obtain

$$\begin{aligned} \int_{B(X, \frac{r}{4})} |\nabla u| |v| dY &\leq \frac{C}{r} \left(\int_{B(X, 0.5r)} |u|^2 dY \right)^{1/2} \left(\int_{B(X, 0.5r)} |v|^2 dY \right)^{1/2} \\ &\leq \frac{C}{r} \sup_{B(X, 0.5r)} u \int_{B(X, 0.5r)} |v| dY \leq \frac{C}{r} \int_{B(X, 0.5r)} |u| |v| dY, \end{aligned}$$

where we used the Harnack's inequality in the last step. The desired estimate now follows by covering $B(x_0, R) \times (r, 2r)$ with balls of radius $r/4$. The proof for the case that $v \geq 0$ is similar. $\square$

The next theorem concerns the uniqueness in the L^p Dirichlet problem in $\mathbb{R}_+^{d+1}$.

Theorem 4.5. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ with coefficients satisfying (1.1)-(1.2). Suppose that $d\omega^Z/dx = K(\cdot, Z) \in L^q(\mathbb{R}^d)$ for all $Z \in \mathbb{R}_+^{d+1}$ and some $q > 1$. Let u be a weak solution to $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ such that $(u)^* \in L^p(\mathbb{R}^d)$, where $p = q'$. Assume that $u(x, t) \rightarrow 0$ as $t \rightarrow 0^+$ for a.e. $x \in \mathbb{R}^d$. Then $u \equiv 0$ in $\mathbb{R}_+^{d+1}$.*

Proof. Fix $Z \in \mathbb{R}_+^{d+1}$. Let $G(X) = G(X, Z)$ be the Green's function with pole at Z . Choose $\varphi(x) \in C_0^\infty(B(0, \ell/2))$ and $\psi(t) \in C_0^\infty(1/(2\ell), 2\ell)$ such that $\varphi = 1$ in $B(0, \ell/4)$, $|\nabla \varphi| \leq C/\ell$ and $\psi = 1$ on $(1/\ell, \ell)$, $|\nabla \psi| \leq C\ell$ on $(1/(2\ell), 1/\ell)$, $|\nabla \psi| \leq C/\ell$ on $(\ell, 2\ell)$. Then

$$\begin{aligned} u(Z) &= \int_{\mathbb{R}_+^{d+1}} a_{ij} \frac{\partial G}{\partial x_j} \cdot \frac{\partial}{\partial x_i} (u\varphi\psi) dY \\ &= \int_{\mathbb{R}_+^{d+1}} a_{ij} \frac{\partial G}{\partial x_j} u \cdot \frac{\partial}{\partial x_i} (\varphi\psi) dY - \int_{\mathbb{R}_+^{d+1}} G \cdot a_{ij} \frac{\partial u}{\partial x_i} \cdot \frac{\partial}{\partial x_j} (\varphi\psi) dY. \end{aligned}$$

This gives

$$\begin{aligned}
|u(Z)| &\leq C\ell \int_{\frac{1}{2\ell}}^{\frac{1}{\ell}} \int_{|y| < \frac{\ell}{2}} (|\nabla G||u| + |G||\nabla u|) dY \\
&\quad + \frac{C}{\ell} \int_{\ell}^{2\ell} \int_{|y| < \frac{\ell}{2}} (|\nabla G||u| + |G||\nabla u|) dY \\
&\quad + \frac{C}{\ell} \int_{\frac{1}{\ell}}^{\ell} \int_{\frac{\ell}{4} < |y| < \frac{\ell}{2}} (|\nabla G||u| + |G||\nabla u|) dY \\
&\leq C\ell^2 \int_{\frac{1}{3\ell}}^{\frac{2}{\ell}} \int_{|y| < \ell} |G||u| dY + \frac{C}{\ell^2} \int_{\frac{\ell}{2}}^{3\ell} \int_{|y| < \ell} |G||u| dY \\
&\quad + \frac{C}{\ell} \int_0^{2\ell} \int_{\frac{\ell}{8} < |y| < \ell} |G||u| \frac{dy dt}{t} \\
&= I_1 + I_2 + I_3,
\end{aligned} \tag{4.6}$$

where we used Lemma 4.4 for the second inequality.

To estimate I_2 , we note that $|G(Y)| \leq C(Z)/\ell^{d-1+\alpha}$ for $Y \in B(0, \ell) \times (\ell/2, 3\ell)$, where $\alpha > 0$. This leads to

$$I_2 \leq \frac{C}{\ell^{d+\alpha}} \int_{B(0, 3\ell)} (u)^* dy \leq C\ell^{-\alpha-\frac{d}{p}} \|(u)^*\|_p \rightarrow 0 \quad \text{as } \ell \rightarrow \infty. \tag{4.7}$$

Next we observe that

$$I_1 \leq C \int_{|y| < \ell} M(K(\cdot, Z)) \mathcal{M}_{2/\ell}(u) dy, \tag{4.8}$$

where M denotes the Hardy-Littlewood maximal operator on $\mathbb{R}^d$ and

$$\mathcal{M}_r(u)(x) = \sup \{u(x, t) : 0 < t < r\}. \tag{4.9}$$

Since $K(\cdot, Z) \in L^{p'}(\mathbb{R}^d)$, we have $I_1 \leq C \|\mathcal{M}_{2/\ell}(u)\|_p$. By the assumption, $\mathcal{M}_{2/\ell}(u)(x) \rightarrow 0$ for a.e. $x \in \mathbb{R}^d$ as $\ell \rightarrow \infty$. Since $\mathcal{M}_{2/\ell}(u) \leq (u)^* \in L^p(\mathbb{R}^d)$, we obtain $\|\mathcal{M}_{2/\ell}(u)\|_p \rightarrow 0$ as $\ell \rightarrow \infty$.

Finally, note that as $\ell \rightarrow \infty$,

$$I_3 \leq C \int_{\frac{\ell}{4} < |y| < \ell} M(K(\cdot, Z))(u)^* dy \leq C \left\{ \int_{|y| > \frac{\ell}{4}} |(u)^*|^p dy \right\}^{1/p} \rightarrow 0.$$

Thus we have proved that $I_1 + I_2 + I_3 \rightarrow 0$ as $\ell \rightarrow \infty$. Hence $u(Z) = 0$ for any $Z \in \mathbb{R}_+^{d+1}$. $\square$

Theorem 4.1 is a direct consequence of Lemma 4.3 and the following theorem.

Theorem 4.6. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ with coefficients satisfying (1.1)-(1.2). Suppose that the L^p Dirichlet problem for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable for some $1 < p < \infty$. Then for any $f \in L^p(\mathbb{R}^d)$, there exists a unique weak solution in $\mathbb{R}_+^{d+1}$ such that $(u)^* \in L^p(\mathbb{R}^d)$ and $u = f$ n.t. on $\mathbb{R}^d$. Moreover the solution satisfies the estimate $\|(u)^*\|_p \leq C\|f\|_p$.*

Proof. Let $f \in C_c(\mathbb{R}^d)$ and $u(X) = \int_{\mathbb{R}^d} f d\omega^X$ be the solution to the classical Dirichlet problem with data f . Since

$$|u(x, t)| \leq \frac{C}{t^d} \int_{B(x, t)} |(u)^*| dy \leq Ct^{-\frac{d}{p}} \|(u)^*\|_p \leq Ct^{-\frac{d}{p}} \|f\|_p,$$

it follows that ω^X is absolutely continuous with respect to dx on $\mathbb{R}^d$ and $K(\cdot, X) = d\omega^X/dx \in L^{p'}(\mathbb{R}^d)$. In view of Theorem 4.5, this gives the uniqueness in Theorem 4.6.

To establish the existence, we let $f \in L^p(\mathbb{R}^d)$ and choose $f_k \in C_c(\mathbb{R}^d)$ so that $f_k \rightarrow f$ in $L^p(\mathbb{R}^d)$. Let u_k be the solution to the classical Dirichlet problem in $\mathbb{R}_+^{d+1}$ with data f_k . It follows from Lemma 4.3 that $\|(u_k)^*\|_p \leq C\|f_k\|_p$ and $\|(u_\ell - u_k)^*\|_p \leq C\|f_\ell - f_k\|_p$. This implies that u_k converges to a weak solution u , uniformly on any compact subset of $\mathbb{R}_+^{d+1}$. Standard limiting arguments show that $\|(u)^*\|_p \leq C\|f\|_p$ and $u = f$ n.t. on $\mathbb{R}^d$. $\square$

5 L^2 estimates for the regularity problem

Definition 5.1. Let $1 < p < \infty$. We say that the L^p regularity problem $(R)_p$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable, if for any $f \in C_0^1(\mathbb{R}^d)$, the unique solution to the classical Dirichlet problem with boundary data f satisfies the estimate $\|N(\nabla u)\|_p \leq C\|\nabla_x f\|_p$.

In this section we study the L^2 regularity problem in $\mathbb{R}_+^{d+1}$ under the condition that the coefficient matrix is periodic in the t direction. As in the case of the Dirichlet problem, we need a local solvability condition: for any $x_0 \in \mathbb{R}^d$ and $0 < r \leq 1$,

$$\int_{B(x_0, r)} |N_{\frac{r}{2}}(\nabla u)|^2 dx \leq C_1 \left\{ \int_{B(x_0, 2r)} |\nabla_x f|^2 dx + \frac{1}{r} \iint_{T(x_0, 2r)} |\nabla u|^2 dx dt \right\}, \quad (5.1)$$

whenever $u \in W^{1,2}(T(x_0, 4r))$ is a weak solution to $\mathcal{L}(u) = 0$ in $T(x_0, 4r)$ such that $u \in C(\overline{T(x_0, 4r)})$ and $u = f \in C^1(B(x_0, 4r))$ on $B(x_0, 4r) \times \{0\}$. We will also assume that the coefficient matrix $A(X)$ is sufficiently smooth. However, constants C in all estimates will depend only on d, μ and C_1 in (5.1).

We begin by observing that since

$$|u(x, t) - u(x, 0)| \leq Ct N_{2t}(\nabla u)(x) \quad (5.2)$$

(see [21], pp.461-462), condition (5.1) implies the local condition (4.1) for the Dirichlet problem. It also follows from (5.1) that

$$\int_{B(x_0, 1)} |N_4(\nabla u)|^2 dx \leq C \left\{ \int_{B(x_0, 2)} |\nabla_x f|^2 dx + \iint_{T(x_0, 6)} |\nabla u|^2 dx dt \right\}. \quad (5.3)$$

Furthermore, if $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$, $u \in C(\mathbb{R}^d \times [0, \infty))$ and $u = f \in C_0^1(\mathbb{R}^d)$ on $\mathbb{R}^d$, we may integrate both sides of (5.3) with respect to x_0 over $\mathbb{R}^d$ to obtain

$$\int_{\mathbb{R}^d} |N_4(\nabla u)|^2 dx \leq C \left\{ \int_{\mathbb{R}^d} |\nabla_x f|^2 dx + \int_0^6 \int_{\mathbb{R}^d} |\nabla u|^2 dx dt \right\}. \quad (5.4)$$

Theorem 5.2. Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ with smooth coefficients satisfying conditions (1.1), (1.2), (1.3). Also assume that $\mathcal{L}$ satisfies the condition (5.1). Then the L^2 regularity problem for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable. Moreover the solution u satisfies the estimate $\|N(\nabla u)\|_2 \leq C\|\nabla_x f\|_2$, where C depends only on d, μ and the constant C_1 in (5.1).

In view of (3.11), we start with a decay estimate.

Lemma 5.3. Under the same assumption on $\mathcal{L}$ as in Theorem 5.2, we have

$$R \iint_{T(0,10R) \setminus T(0,R)} |\nabla u|^2 dx dt \rightarrow 0 \quad \text{as } R \rightarrow \infty, \quad (5.5)$$

where u is a classical solution to $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ such that $u = f \in C_c(\mathbb{R}^d)$ on $\mathbb{R}^d$.

Proof. By Cacciopoli's inequality,

$$\begin{aligned} R \iint_{T(0,10R) \setminus T(0,R)} |\nabla u|^2 dx dt &\leq \frac{C}{R} \iint_{T(0,11R) \setminus T(0,R/2)} |u|^2 dx dt \\ &\leq C \int_{|x| \geq \frac{R}{100}} |(u)^*|^2 dx \end{aligned} \quad (5.6)$$

for R sufficiently large. Since $(u)^* \in L^2(\mathbb{R}^d)$, the right-hand side of (5.6) goes to zero as $R \rightarrow \infty$. $\square$

Proof of Theorem 5.2. Let $f \in C_0^1(\mathbb{R}^d)$ and u be the solution to the classical Dirichlet problem with data f . It follows from Remark 2.2 and (5.4) that

$$\begin{aligned} \int_{\mathbb{R}^d} |N(\nabla u)|^2 dx &\leq C \left\{ \int_{\mathbb{R}^d} |\nabla_x f|^2 dx + \int_{\mathbb{R}^d} |N_4(\nabla u)|^2 dx + \int_{\mathbb{R}^d} |(Q(u))^*|^2 dx \right\} \\ &\leq C \left\{ \int_{\mathbb{R}^d} |\nabla_x f|^2 dx + \int_0^6 \int_{\mathbb{R}^d} |\nabla u|^2 dx dt + \int_{\mathbb{R}^d} |(Q(u))^*|^2 dx \right\}. \end{aligned} \quad (5.7)$$

To estimate the integral of $|\nabla u|^2$ over $\mathbb{R}^d \times (0,6)$, we let $R \rightarrow \infty$ in (3.11). In view of (5.5), we obtain

$$\int_0^6 \int_{\mathbb{R}^d} |\nabla u|^2 dx dt \leq C \int_{\mathbb{R}^d} |\nabla_x f|^2 dx. \quad (5.8)$$

Finally, to handle the term with $Q(u)$ in (5.7), we observe that u is also the unique solution to the L^2 Dirichlet problem for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ with boundary data f . This implies that $(Q(u))^* \in L^2(\mathbb{R}^d)$ and by Theorem 4.1,

$$\begin{aligned} \int_{\mathbb{R}^d} |(Q(u))^*|^2 dx &\leq C \int_{\mathbb{R}^d} |Q(u)|^2 dx \leq C \int_0^1 \int_{\mathbb{R}^d} |\nabla u|^2 dx dt \\ &\leq C \int_{\mathbb{R}^d} |\nabla_x f|^2 dx, \end{aligned} \quad (5.9)$$

where we used (5.8) in the last step. The desired estimate $\|N(\nabla u)\|_2 \leq C\|\nabla_x f\|_2$ now follows from (5.7), (5.8) and (5.9). $\square$

We omit the proof of the next lemma and refer the reader to [21] for an analogous result (Theorem 3.1, pp.461-462) in the case of the unit ball.

Lemma 5.4. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ be an elliptic operator satisfying (1.1)-(1.2). Suppose that $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ and $N_r(\nabla u) \in L^p(\mathbb{R}^d)$ for some $1 < p < \infty$ and $r > 0$. Then u converges n.t. to f on $\mathbb{R}^d$ and $\nabla_x f \in L^p(\mathbb{R}^d)$. Moreover, let*

$$V_i(x, t) = \frac{1}{t|B(x, t)|} \int_{\frac{t}{2}}^{\frac{3t}{2}} \int_{B(x, t)} \frac{\partial u}{\partial y_i} dy ds$$

for $i = 1, \dots, d$, then $V_i(x, t)$ converges weakly to $\frac{\partial f}{\partial x_i}$ in $L^p(\mathbb{R}^d)$ as $t \rightarrow 0$. In particular, $\|\nabla_x f\|_p \leq \|N(\nabla u)\|_p$.

The next theorem concerns the uniqueness of solutions to the L^p regularity problem.

Theorem 5.5. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ with coefficients satisfying (1.1)-(1.2). Let u be a weak solution to $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ such that $N(\nabla u) \in L^p(\mathbb{R}^d)$ for some $1 \leq p < \frac{d}{1-\alpha}$, where $\alpha > 0$ is given by (2.2). Then $u \equiv 0$ in $\mathbb{R}_+^{d+1}$, if $u(x, t) \rightarrow 0$ as $t \rightarrow 0^+$ for a.e. $x \in \mathbb{R}^d$.*

Proof. The proof begins in the same way as that of Theorem 4.5. As such, this leads to $|u(Z)| \leq I_1 + I_2 + I_3$, where I_1 , I_2 and I_3 are defined in (4.6). To estimate I_1 , we use the estimate (2.2) and the observation that $|u(x, t)| \leq CtN(\nabla u)(x)$ for a.e. $x \in \mathbb{R}^d$. This yields that

$$I_1 = C\ell^2 \int_{\frac{1}{2\ell}}^{\frac{2}{\ell}} \int_{|y| < \ell} |G||u| dY \leq \frac{C}{\ell^\alpha} \int_{\mathbb{R}^d} \frac{N(\nabla u)(y) dy}{|(y, 0) - Z|^{d-1+\alpha}} \leq C\ell^{-\alpha} \|N(\nabla u)\|_p \rightarrow 0$$

as $\ell \rightarrow \infty$, where we used the assumption $p < \frac{d}{1-\alpha}$. Similarly,

$$I_2 = \frac{C}{\ell^2} \int_{\frac{\ell}{2}}^{3\ell} \int_{|y| < \ell} |G||u| dy dt \leq \frac{C}{\ell^{d-1+\alpha}} \int_{|y| < \ell} N(\nabla u) dy \leq C\ell^{1-\alpha-\frac{d}{p}} \|N(\nabla u)\|_p$$

which also tends to zero as $\ell \rightarrow \infty$, since $p < \frac{d}{1-\alpha}$. Finally, note that

$$I_3 = \frac{C}{\ell} \int_0^{2\ell} \int_{\frac{\ell}{4} < |y| < 2\ell} |G||u| \frac{dy dt}{t} \leq \frac{C}{\ell^{d-1+\alpha}} \int_{|y| < 2\ell} |N(\nabla u)| dy \rightarrow 0,$$

as $\ell \rightarrow \infty$. Thus we have proved that $I_1 + I_2 + I_3 \rightarrow 0$ as $\ell \rightarrow \infty$. Hence $u(Z) = 0$ for any $Z \in \mathbb{R}_+^{d+1}$. $\square$

Recall that $f \in \dot{W}^{1,p}(\mathbb{R}^d)$ if $f \in W_{loc}^{1,p}(\mathbb{R}^d)$ and $\nabla_x f \in L^p(\mathbb{R}^d)$. The following theorem provides the existence of solutions with boundary data in $\dot{W}^{1,p}(\mathbb{R}^d)$.

Theorem 5.6. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ with coefficients satisfying (1.1)-(1.2). Suppose that the L^p regularity problem for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable. Then for any $f \in \dot{W}^{1,p}(\mathbb{R}^d)$, there exists a solution u to $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ such that $\|N(\nabla u)\|_p \leq C\|f\|_p$ and $u = f$ n.t. on $\mathbb{R}^d$.*

Proof. Let $f \in \dot{W}^{1,p}(\mathbb{R}^d)$ and choose $f_k \in C_0^1(\mathbb{R}^d)$ such that $\nabla_x f_k \rightarrow \nabla_x f$ in $L^p(\mathbb{R}^d)$. Let v_k be the solution to the classical Dirichlet problem in $\mathbb{R}_+^{d+1}$ with boundary data f_k . By

assumption, we have $\|N(\nabla v_k)\|_p \leq C\|\nabla_x f_k\|_p$ and $\|N(\nabla v_k - \nabla v_\ell)\|_p \leq C\|\nabla_x(f_k - f_\ell)\|_p$. It follows that for any $m > 1$, $\nabla(v_k - v_\ell) \rightarrow 0$ in $L^2(\Omega_m)$ as $k, \ell \rightarrow \infty$, where

$$\Omega_m = \left\{ (x, t) \in \mathbb{R}_+^{d+1} : |x| < 3m \text{ and } \frac{1}{3m} < t < 3m \right\}.$$

Thus there exists $u_m \in W^{1,2}(\Omega_m)$ such that $v_k - \alpha(k, m) \rightarrow u_m$ in $W^{1,2}(\Omega_m)$ as $k \rightarrow \infty$, where $\alpha(k, m)$ is the average of v_k over Ω_m . Clearly, $\mathcal{L}(u_m) = 0$ in Ω_m . By a limiting argument, we also have

$$\|\tilde{N}_m(\nabla u_m)\|_{L^p(B(0,m))} \leq C\|\nabla_x f\|_p, \quad (5.10)$$

where the nontangential maximal function $\tilde{N}_m(\nabla u)$ is defined in a manner similar to N_m , but the variable t in (2.9) is restricted to $(1/m) < t < m$.

Next, since $u_{m+1} - u_m$ is constant in Ω_m , one may define a function $u \in W_{loc}^{1,p}(\mathbb{R}_+^{d+1})$ such that $u - u_m$ is constant in Ω_m for any $m \geq 1$. It follows that u is a weak solution in $\mathbb{R}_+^{d+1}$. Moreover, in view of (5.10), we have

$$\|\tilde{N}_m(\nabla u)\|_{L^p(B(0,m))} \leq C\|\nabla_x f\|_p. \quad (5.11)$$

Letting $m \rightarrow \infty$ in (5.11) gives $\|N(\nabla u)\|_p \leq C\|\nabla_x f\|_p$. Similar argument also gives

$$\|N(\nabla u - \nabla v_k)\|_p \leq C\|\nabla_x(f - f_k)\|_p. \quad (5.12)$$

Finally we note that by Lemma 5.4, u converges n.t. to $h \in \dot{W}^{1,p}(\mathbb{R}^d)$ and the average of $\frac{\partial u}{\partial x_i}$ over $B(x, t) \times (t/2, 3t/2)$ converges weakly to $\frac{\partial h}{\partial x_i}$ in $L^p(\mathbb{R}^d)$ as $t \rightarrow 0$. This, together with the weak convergence of the average of $\frac{\partial v_k}{\partial x_i}$ to $\frac{\partial f_k}{\partial x_i}$ and estimate (5.12), implies that $\frac{\partial h}{\partial x_i} = \frac{\partial f}{\partial x_i}$ on $\mathbb{R}^d$ for $i = 1, \dots, d$. It follows that $f - h$ is constant. Thus, by subtracting a constant, we obtain $u = f$ n.t. on $\mathbb{R}^d$. $\square$

We conclude this section with two more theorems on the consequences of the solvability of the regularity problem.

Theorem 5.7. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ with coefficients satisfying (1.1)-(1.2). Let $1 < p, q < \infty$ and $\frac{1}{p} + \frac{1}{q} = 1$. Then for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$, the solvability of the L^p regularity problem implies the solvability of the L^q Dirichlet problem.*

Theorem 5.8. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ with coefficients satisfying (1.1)-(1.2). Suppose that the L^2 regularity problem for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable. Let $u \in W^{1,2}(T(x_0, 8r))$ be a weak solution in $T(x_0, 8r)$ such that $N_r(\nabla u) \in L^2(B(x_0, 4r))$ and $u = f$ on $B(x_0, 4r)$. Then*

$$\int_{B(x_0, r)} |N_{\frac{r}{2}}(\nabla u)|^2 dx \leq C \int_{B(x_0, 2r)} |\nabla_x f|^2 dx + \frac{C}{r} \int_{T(x_0, 2r)} |\nabla u|^2 dX. \quad (5.13)$$

We omit the proof of both theorems and refer the reader to Theorems 5.4 and 5.19 in [21], where the analogous results were proved in the case of the unit ball. We point out that Theorem 5.19 in [21] was stated for solutions in the whole domain. However an inspection of its proof, which extends readily to the case of $\mathbb{R}_+^{d+1}$, shows that the conclusion holds for local solutions. Theorem 5.8 shows that the local solvability condition (5.1) is necessary.

6 L^p estimates for the regularity problem

In this section we study the L^p regularity problem in $\mathbb{R}_+^{d+1}$ for $1 < p < 2 + \delta$.

Theorem 6.1. *Let $\mathcal{L} = -\operatorname{div}(A\nabla u)$ with coefficients satisfying (1.1)-(1.2). Suppose that $(R)_2$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable. Then $(R)_p$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable for $1 < p < 2 + \delta$.*

In the case of the unit ball, Kenig and Pipher [21] proved that the solvability of $(R)_q$ implies the solvability of $(R)_p$ for $1 < p < q + \delta$. Although it is possible to extend their results to the case of the upper half-space and thus obtain the L^p solvability for $1 < p < 2 + \delta$ from the L^2 solvability, we will provide a more direct proof.

For the range $1 < p < 2$, we follow the approach used in [13] (also see [21]) by proving $\|N(\nabla u)\|_1 \leq C$ for solutions of the L^2 regularity problem with H_{at}^1 data. Our approach to the range $2 < p < 2 + \delta$, which is based on a real variable argument developed by Shen [25, 26], is different from that used in [13, 21]. The rather general approach reduces the problem to certain weak reverse Hölder estimates. In particular, our argument shows that if $(R)_2$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable, then the solvability of $(D)_q$ for some $q < 2$ implies the solvability of $(R)_p$, where $\frac{1}{p} + \frac{1}{q} = 1$.

We start with a comparison principle.

Lemma 6.2. *Let $u, v \in C(\overline{T(x_0, 3r)})$ be two weak solutions in $T(x_0, 3r)$ such that $u(x, 0) = v(x, 0) = 0$ on $B(x_0, 3r)$. Suppose that v is nonnegative in $T(x_0, 3r)$. Then*

$$|u(x, t)| \leq C \cdot \frac{v(x, t)}{v(x_0, r)} \left(\frac{1}{r^{d+1}} \iint_{T(x_0, 3r)} |u(y, s)|^2 dy ds \right)^{1/2}$$

for any $(x, t) \in T(x_0, r)$, where C depends only on d and μ .

Proof. See Lemma 2.5 in [27]. □

The following is a localization result similar to Theorem 5.8. Note that here we assume the solvability of $(D)_2$ instead of the solvability of $(R)_2$.

Theorem 6.3. *Let $\mathcal{L}$ be an elliptic operator with coefficients satisfying (1.1)-(1.2). Assume that the L^2 Dirichlet problem for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable. Then*

$$\int_{B(x_0, r)} |N_r(\nabla u)|^2 dx \leq \frac{C}{r^3} \iint_{T(x_0, 4r)} |u(x, t)|^2 dx dt, \quad (6.1)$$

where $u \in C(\overline{T(x_0, 6r)})$ is a weak solution of $\mathcal{L}(u) = 0$ in $T(x_0, 6r)$ and $u(x, 0) = 0$ on $B(x_0, 6r)$.

Proof. Let $v(x, t)$ be the Green's function for $\mathcal{L}$ on $\mathbb{R}_+^{d+1}$ with pole at $X_0 = (x_0, 10r)$. Fix $x \in B(x_0, r_0)$ and $0 < t < \frac{r}{2}$. By Cacciopoli's inequality,

$$\begin{aligned} \frac{1}{t|B(x, t)|} \int_{\frac{t}{2}}^{\frac{3t}{2}} \int_{B(x, t)} |\nabla u|^2 dy ds &\leq \frac{C}{t^3|B(x, t)|} \int_{\frac{t}{4}}^{2t} \int_{B(x, 2t)} |u(y, s)|^2 dy ds \\ &\leq \frac{C}{t^3|B(x_0, t)|} \int_{\frac{t}{4}}^{2t} \int_{B(x, 2t)} \left| \frac{v(y, s)}{v(x_0, r)} \right|^2 dy ds \cdot \frac{1}{r^{d+1}} \iint_{T(x_0, 4r)} |u(y, s)|^2 dy ds \\ &\leq Cr^{2d-2} \left(\frac{v(x, t)}{t} \right)^2 \cdot \frac{1}{r^{d+1}} \iint_{T(x_0, 4r)} |u(y, s)|^2 dy ds, \end{aligned}$$

where we have used Lemma 6.2 and $v(x_0, r) \approx r^{1-d}$. Let ω denote the $\mathcal{L}$ -harmonic measure for $\mathbb{R}_+^{d+1}$, evaluated at $(x_0, 10r)$. Since

$$\frac{v(x, t)}{t} \approx \frac{\omega(B(x, t))}{|B(x, t)|} \leq CM(\chi_{B(x_0, 2r)} \frac{d\omega}{dx})(x)$$

where M denotes the Hardy-Littlewood maximal operator on $\mathbb{R}^d$, we obtain

$$N_r(\nabla u)(x) \leq Cr^{d-1} M(\chi_{B(x_0, 2r)} \frac{d\omega}{dx})(x) \cdot \left(\frac{1}{r^{d+1}} \iint_{T(x_0, 4r)} |u(y, s)|^2 dy ds \right)^{1/2}. \quad (6.2)$$

Since the L^2 Dirichlet problem for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable, the reverse Hölder inequality (4.3) for $K = \frac{d\omega}{dx}$ holds on $B(x_0, 2r)$. The desired estimate now follows by the L^2 boundedness of M and (4.3). $\square$

Remark 6.4. Suppose that $(D)_p$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable for some $p < 2$. Then the reverse Hölder inequality (4.5) holds for $q = p'$. It follows from (6.2) and (4.5) that

$$\begin{aligned} \left(\frac{1}{r^d} \int_{B(x_0, r)} |N_r(\nabla u)|^q dx \right)^{1/q} &\leq \frac{C}{r} \left(\frac{1}{r^{d+1}} \iint_{T(x_0, 4r)} |u(y, s)|^2 dy ds \right)^{1/2} \\ &\leq C \left(\frac{1}{r^{d+1}} \iint_{T(x_0, 4r)} |\nabla u(y, s)|^2 dy ds \right)^{1/2} \\ &\leq C \left(\frac{1}{r^d} \int_{B(x_0, 6r)} |N(\nabla u)|^2 dx \right)^{1/2}. \end{aligned} \quad (6.3)$$

A simple geometric observation shows that for $x \in B(x_0, r)$,

$$N(\nabla u)(x) \leq N_r(\nabla u)(x) + \frac{C}{r^d} \int_{B(x_0, 6r)} |N(\nabla u)| dx. \quad (6.4)$$

In view of (6.3), this gives

$$\left(\frac{1}{r^d} \int_{B(x_0, r)} |N(\nabla u)|^q dx \right)^{1/q} \leq C \left(\frac{1}{r^d} \int_{B(x_0, 6r)} |N(\nabla u)|^2 dx \right)^{1/2}. \quad (6.5)$$

We will need this weak reverse Hölder estimate to treat the case $2 < p < 2 + \delta$ for the regularity problem.

Theorem 6.5. *Under the same conditions on $\mathcal{L}$ as in Theorem 6.1, the L^p regularity problem in $\mathbb{R}_+^{d+1}$ is solvable for $1 < p < 2$.*

Proof. We follow the approach in [13] and prove that $\|N(\nabla u)\|_1 \leq C$ for solutions of the L^2 regularity problem with H_{at}^1 data. To this end, let f be a Lipschitz function such that $\text{supp}(f) \subset B(x_0, r)$ and $\|\nabla_x f\|_\infty \leq Cr^{-d}$ for some $r > 0$ and $x_0 \in \mathbb{R}^d$. Let u be the solution of the L^2 regularity problem with data f . By the L^2 estimate $\|N(\nabla u)\|_2 \leq C\|\nabla_x f\|_2$, one has $\|N(\nabla u)\|_{L^1(B(x_0, Cr))} \leq C$. Thus it suffices to show that if $R \geq r$ and $B(y_0, 10R) \cap B(x_0, r) = \emptyset$,

$$\int_{B(y_0, R)} |N(\nabla u)| dx \leq C \left(\frac{r}{R}\right)^\alpha \quad (6.6)$$

for some $\alpha > 0$. This follows from Theorem 6.3 and the pointwise estimate

$$\begin{aligned} |u(X)| &\leq \|f\|_\infty \omega^X(B(x_0, r)) \leq Cr^{1-d} \omega^X(B(x_0, r)) \\ &\leq CG(X, X_0) \leq \frac{Cr^\alpha}{|X - X_0|^{d-1+\alpha}}, \end{aligned}$$

where $X_0 = (x_0, r)$. We omit the details. $\square$

If Ω is a bounded Lipschitz domain, it was proved in [27] that the solvability of $(R)_p$ for $\mathcal{L}(u) = 0$ in Ω is equivalent to the solvability of $(D)_{p'}$ for $\mathcal{L}(u) = 0$ in Ω , provided that $(R)_{p_0}$ for $\mathcal{L}(u) = 0$ in Ω is solvable for some $p_0 > 1$. We extend this result to the case $\Omega = \mathbb{R}_+^{d+1}$.

Theorem 6.6. *Let $\mathcal{L} = -\text{div}(A\nabla)$ with coefficients satisfying (1.1)-(1.2). Suppose that $(R)_2$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable. Let $p > 2$ and $q = p'$. Then $(R)_p$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable if and only if $(D)_q$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable.*

The proof of Theorem 6.6 relies on a real variable argument which may be formulated as follows.

Theorem 6.7. *Let $F \in L^2(\mathbb{R}^d)$ and $g \in L^p(\mathbb{R}^d)$ for some $2 < p < q$. Also assume that g has compact support. Suppose that for each ball B in $\mathbb{R}^d$, there exist two functions F_B and R_B such that $|F| \leq |F_B| + |R_B|$ on $2B$, and*

$$\begin{aligned} \frac{1}{|2B|} \int_{2B} |F_B|^2 dx &\leq E_1 \sup_{B' \supset B} \frac{1}{|B'|} \int_{B'} |g|^2 dx, \\ \left\{ \frac{1}{|2B|} \int_{2B} |R_B|^q dx \right\}^{1/q} &\leq E_2 \left\{ \left(\frac{1}{|\beta B|} \int_{\beta B} |F|^2 dx \right)^{1/2} + \sup_{B' \supset B} \left(\frac{1}{|B'|} \int_{B'} |g|^2 dx \right)^{1/2} \right\}, \end{aligned}$$

where $E_1, E_2 > 0$ and $\beta > 2$. Then $F \in L^p(\mathbb{R}^d)$ and $\|F\|_p \leq C\|g\|_p$, where C depends only on d, p, q, E_1, E_2 and β .

Proof. This is a simple consequence of Theorem 3.2 in [26]. Indeed, it follows from the theorem that

$$\left\{ \int_{B(0, r)} |F|^p dx \right\}^{1/p} \leq C|B(0, r)|^{\frac{1}{p}-\frac{1}{2}} \left\{ \int_{B(0, Cr)} |F|^2 dx \right\}^{1/2} + C \left\{ \int_{B(0, Cr)} |g|^p dx \right\}^{1/p}. \quad (6.7)$$

The estimate $\|F\|_p \leq C\|g\|_p$ follows by letting $r \rightarrow \infty$ in (6.7). $\square$

Proof of Theorem 6.6. In view of Theorem 5.7, we only need to show that the solvability of $(D)_q$ implies the solvability of $(R)_p$. More precisely, it suffices to show that $\|N(\nabla u)\|_p \leq C\|\nabla_x f\|_p$, if u is the solution of the classical Dirichlet problem with data $f \in C_0^1(\mathbb{R}^d)$. This will be done by applying Theorem 6.7 to the functions $F = N(\nabla u)$ and $g = |\nabla_x f|$.

To this end, we fix a ball $B = B(x_0, r)$ in $\mathbb{R}^d$. Choose $\varphi \in C_0^1(B(x_0, 8r))$ such that $\varphi = 1$ in $B(x_0, 7r)$ and $|\nabla \varphi| \leq Cr^{-1}$. Let λ be the average of f over $B(x_0, 8r)$. Write $u - \lambda = v + w$, where v is the solution of the L^2 regularity problem with boundary data $(f - \lambda)\varphi$.

We now let $F_B = N(\nabla v)$ and $R_B = N(\nabla w)$. Clearly, $F = N(\nabla u) \leq F_B + R_B$. By the L^2 regularity estimate,

$$\begin{aligned} \frac{1}{|2B|} \int_{2B} |F_B|^2 dx &\leq \frac{1}{|B|} \int_{\mathbb{R}^d} |N(\nabla u)|^2 dx \leq \frac{C}{|B|} \int_{\mathbb{R}^d} |\nabla_x((f - \lambda)\varphi)|^2 dx \\ &\leq \frac{C}{|B|} \int_{B(x_0, 8r)} |\nabla_x f|^2 dx \leq C \sup_{B' \supset B} \frac{1}{|B'|} \int_{B'} |g|^2 dx, \end{aligned}$$

where we have used the Poincaré inequality. This gives the estimate needed for F_B . To verify the condition on $R_B = N(\nabla w)$, we note that w is the solution of the L^2 regularity problem with data $(f - \lambda)(1 - \varphi)$. Consequently, $w = 0$ on $B(x_0, 7r)$ and we may use the weak reverse Hölder estimate (6.5). This leads to

$$\begin{aligned} \left\{ \frac{1}{|2B|} \int_{2B} |R_B|^q dx \right\}^{1/q} &= \left\{ \frac{1}{|2B|} \int_{2B} |N(\nabla w)|^q dx \right\}^{1/q} \\ &\leq C \left\{ \frac{1}{|B|} \int_{B(x_0, 12r)} |N(\nabla w)|^2 dx \right\}^{1/2} \\ &\leq C \left\{ \frac{1}{|B|} \int_{B(x_0, 12r)} |N(\nabla u)|^2 dx \right\}^{1/2} + C \left\{ \frac{1}{|B|} \int_{B(x_0, 12r)} |N(\nabla v)|^2 dx \right\}^{1/2} \\ &\leq C \left\{ \frac{1}{|B(x_0, 12r)|} \int_{B(x_0, 12r)} |F|^2 dx \right\}^{1/2} + C \sup_{B' \supset B} \left\{ \frac{1}{|B'|} \int_{B'} |g|^2 dx \right\}^{1/2}, \end{aligned}$$

which gives the estimate needed for R_B . Therefore, by Theorem 6.7, we obtain $\|N(\nabla u)\|_p \leq C\|\nabla_x f\|_p$ for $2 < p < q$. Finally, since the weak reverse Hölder estimate (6.5) is self-improving, the argument above in fact gives $\|N(\nabla u)\|_p \leq C\|\nabla_x f\|_p$ for $2 < p < q + \delta$. This finishes the proof. $\square$

7 A Neumann function on $\mathbb{R}_+^{d+1}$

Throughout this section we will assume that $\mathcal{L} = -\operatorname{div}(A\nabla)$ with coefficients satisfying (1.1)-(1.2).

Definition 7.1. Let $g \in L_{loc}^1(\mathbb{R}^d)$. We call $u \in W_{loc}^{1,2}(\mathbb{R}_+^{d+1})$ a weak solution to the Neumann problem,

$$\mathcal{L}(u) = 0 \quad \text{in } \Omega = \mathbb{R}_+^{d+1} \quad \text{and} \quad \frac{\partial u}{\partial \nu} = g \quad \text{on } \partial\Omega, \quad (7.1)$$

if $|\nabla u| \in L^1(T(0, R))$ for any $R > 1$ and

$$\int_{\mathbb{R}_+^{d+1}} a_{ij} \frac{\partial u}{\partial x_j} \frac{\partial \varphi}{\partial x_i} dX = \int_{\mathbb{R}^d} g \varphi dx \quad (7.2)$$

for any $\varphi \in C_0^1(\mathbb{R}^{d+1})$.

To construct weak solutions to (7.1) we introduce a Neumann function. Let $E = (e_{ij})$ be the $(d+1) \times (d+1)$ diagonal matrix with $e_{11} = \dots = e_{dd} = 1$ and $e_{(d+1)(d+1)} = -1$, and

$$\tilde{A} = \tilde{A}(x, t) = \begin{cases} A(x, t) & \text{if } t > 0, \\ EA(x, -t)E & \text{if } t < 0, \end{cases} \quad (7.3)$$

where $A(x, t)$ is the coefficient matrix for $\mathcal{L}$. Let $\tilde{\Gamma} = \tilde{\Gamma}(X, Y)$ denote the fundamental solution for the operator $\tilde{L} = -\operatorname{div}(\tilde{A}\nabla)$ on $\mathbb{R}^{d+1}$ with pole at X . Using the fact that $\tilde{L}(v) = 0$ if $v(x, t) = u(x, -t)$ and $\tilde{L}(u) = 0$, one may show that

$$\tilde{\Gamma}(X^*, Y^*) = \tilde{\Gamma}(X, Y) \quad \text{for any } X, Y \in \mathbb{R}^{d+1}, \quad (7.4)$$

where $X^* = (x, -t)$ for $X = (x, t)$. We now define

$$N(X, Y) = \tilde{\Gamma}(X, Y) + \tilde{\Gamma}(X, Y^*) \quad \text{for } X, Y \in \mathbb{R}_+^{d+1}. \quad (7.5)$$

Clearly, $N(X, Y) = N(Y, X)$. Since $|X - Y| \leq |X - Y^*|$ for $X, Y \in \mathbb{R}_+^{d+1}$, we also have

$$\begin{aligned} |N(X, Y)| &\leq \frac{C}{|X - Y|^{d-1}}, \\ |N(X, Y) - N(Z, Y)| &\leq \frac{C|X - Z|^\alpha}{|X - Y|^{d-1+\alpha}}, \end{aligned} \quad (7.6)$$

for any $X, Y, Z \in \mathbb{R}_+^{d+1}$, where $C > 0$, $\alpha > 0$ depend only on d and μ .

Let $N(X, y) = N(X, (y, 0))$ for $y \in \mathbb{R}^d$. The next lemma shows that $N(X, Y)$ is a Neumann function for the operator $\mathcal{L}$ on $\mathbb{R}_+^{d+1}$.

Lemma 7.2. (a) For $1 \leq p \leq 2$, let $g \in L^p(\mathbb{R}^d)$ and

$$u(X) = \begin{cases} \int_{\mathbb{R}^d} N(X, y)g(y) dy & \text{for } d \geq 3 \text{ or } d = 2, 1 \leq p < 2, \\ \int_{\mathbb{R}^d} (N(X, y) - N(X_0, y))g(y) dy & \text{for } d = 2 \text{ and } p = 2, \end{cases} \quad (7.7)$$

where $X_0 = (0, 1) \in \mathbb{R}_+^{d+1}$. Then u is a weak solution to the Neumann problem on $\mathbb{R}_+^{d+1}$ with data g .

(b) Let $F \in C_c(\mathbb{R}^{d+1})$ and

$$w(X) = \int_{\mathbb{R}_+^{d+1}} N(X, Y)F(Y) dY. \quad (7.8)$$

Then

$$\int_{\mathbb{R}_+^{d+1}} a_{ij} \frac{\partial w}{\partial x_j} \frac{\partial \varphi}{\partial x_i} dX = \int_{\mathbb{R}_+^{d+1}} F \varphi dX \quad (7.9)$$

for any $\varphi \in C_0^1(\mathbb{R}^{d+1})$.

(c) If $g \in L^2(\mathbb{R}^d)$ and has compact support, then the solution u defined by (7.7) belongs to $W^{1,2}(T(0, R))$ for any $R \geq 1$.

Proof. Part (a) follows from

$$\varphi(Y) = 2 \int_{\mathbb{R}_+^{d+1}} a_{ij} \frac{\partial}{\partial x_j} \{ \tilde{\Gamma}(X, Y) \} \frac{\partial \varphi}{\partial x_i} dX \quad (7.10)$$

for $Y = (y, 0)$ and $\varphi \in C_0^1(\mathbb{R}^{d+1})$. To see this, one uses

$$\tilde{\varphi}(Y) = \int_{\mathbb{R}^{d+1}} \tilde{a}_{ij} \frac{\partial}{\partial x_j} \{ \tilde{\Gamma}(X, Y) \} \frac{\partial \tilde{\varphi}}{\partial x_i} dX,$$

where $\tilde{\varphi}$ is the even reflection of φ from $\mathbb{R}_+^{d+1}$ to $\mathbb{R}^{d+1}$, i.e., $\tilde{\varphi}(X) = \varphi(X)$ if $x_{d+1} \geq 0$ and $\tilde{\varphi}(X) = \varphi(X^*)$ if $x_{d+1} < 0$.

To see (b), we write

$$w(X) = \int_{\mathbb{R}^{d+1}} \tilde{\Gamma}(X, Y) \tilde{F}(Y) dY$$

where $\tilde{F}$ is the even reflection of F and deduce (7.9) from

$$\int_{\mathbb{R}^{d+1}} \tilde{a}_{ij} \frac{\partial w}{\partial x_j} \frac{\partial \tilde{\varphi}}{\partial x_i} dX = \int_{\mathbb{R}^{d+1}} \tilde{F} \tilde{\varphi} dX.$$

Part (c) follows by duality from part (b). Let $h \in C_0^1(\mathbb{R}_+^{d+1})$. Note that

$$\begin{aligned} \int_{\mathbb{R}_+^{d+1}} \frac{\partial u}{\partial x_j} h(X) dX &= 2 \int_{\mathbb{R}^d} g(y) dy \int_{\mathbb{R}_+^{d+1}} \frac{\partial \tilde{\Gamma}}{\partial x_j}(X, (y, 0)) h(X) dX \\ &= -2 \int_{\mathbb{R}^d} g(y) v_j(y) dy, \end{aligned}$$

where

$$v_j(y) = \int_{\mathbb{R}_+^{d+1}} \tilde{\Gamma}(X, (y, 0)) \frac{\partial h}{\partial x_j} dX.$$

Using

$$\|v_j\|_{L^2(B(0, R))} \leq C_R \|v_j\|_{W^{1,2}(T(0, R))} \leq C_R \|h\|_{L^2(\mathbb{R}_+^{d+1})},$$

we may conclude by duality that $|\nabla u| \in L^2(T(0, R))$ for any $R > 1$. $\square$

Definition 7.3. Let $1 < p < \infty$. We say the L^p Neumann problem $(N)_p$ for $\mathbb{R}_+^{d+1}$ is solvable if for any $g \in L_c^\infty(\mathbb{R}^d)$, the weak solution $u(X) = \int_{\mathbb{R}^d} N(X, y) f(y) dy$ satisfies $\|N(\nabla u)\|_p \leq C \|g\|_p$.

The solvability of $(N)_p$ yields the existence of weak solutions of the Neumann problem with data in $L^p(\mathbb{R}^d)$.

Theorem 7.4. *Let $1 < p < \infty$. Suppose that $(N)_p$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable. Then for any $g \in L^p(\mathbb{R}^d)$, there exists a weak solution u to the Neumann problem with data g such that $\|N(\nabla u)\|_p \leq C\|g\|_p$.*

Proof. Let $g \in L^p(\mathbb{R}^d)$. Choose $g_k \in C_c(\mathbb{R}^d)$ such that $g_k \rightarrow g$ in $L^p(\mathbb{R}^d)$. Let $u_k(X) = \int_{\mathbb{R}^d} N(X, y)g_k(y)dy$. Then $\|N(\nabla v_k)\|_p \leq C\|g_k\|_p$ and $\|N(\nabla v_k - \nabla v_\ell)\|_p \leq C\|g_k - g_\ell\|_p$. By a limiting argument similar to that in the proof of Theorem 5.6, there exists $u \in W_{loc}^{1,2}(\mathbb{R}_+^{d+1})$ such that $\|N(\nabla u)\|_p \leq C\|g\|_p$ and $\|N(\nabla u_k - \nabla u)\|_p \leq C\|g_k - g\|_p$. Note that the last inequality implies $\nabla u_k \rightarrow \nabla u$ in $L^p(T(0, R))$ for any $R > 1$. It follows that u is a weak solution to the Neumann problem with data g . $\square$

Remark 7.5. Suppose that $(N)_p$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable. It follows from the proof of Theorem 7.4 that if $1 < p \leq 2$, the weak solution given by (7.7) satisfies $\|N(\nabla u)\|_p \leq C\|g\|_p$.

Lemma 7.6. *Let $u \in W_{loc}^{1,2}(T(0, 2R))$. Assume that*

$$\int_{\mathbb{R}_+^{d+1}} a_{ij} \frac{\partial u}{\partial x_j} \cdot \frac{\partial \psi}{\partial x_i} dX = 0 \quad (7.11)$$

for any $\psi \in C_0^1(B(0, 2R) \times (-R, R))$. Then

$$|u(X) - u(Y)| \leq CR \left(\frac{|X - Y|}{R} \right)^\beta \left(\frac{1}{R^{d+1}} \int_{T(0, 2R)} |\nabla u|^2 dX \right)^{1/2} \quad (7.12)$$

for any $X, Y \in T(0, R)$, where $C > 0$ and $\beta \in (0, 1]$ depend only on d and μ .

Proof. Estimate (7.12) follows from the De Giorgi - Nash estimate by a reflection argument. $\square$

The next theorem addresses the question of uniqueness.

Theorem 7.7. *Let $u \in W_{loc}^{1,2}(\mathbb{R}_+^{d+1})$ and $N(\nabla u) \in L^p(\mathbb{R}^d)$ for some $2 \leq p < \frac{d}{1-\beta}$, where β is given by (7.12). Suppose that (7.11) holds for any $\psi \in C_0^1(\mathbb{R}^{d+1})$. Then u is constant in $\mathbb{R}_+^{d+1}$.*

Proof. This follows readily from the estimate (7.12). $\square$

We end this section with a result on the implication of $N(\nabla u) \in L^p(\mathbb{R}^d)$ on the Neumann data.

Theorem 7.8. *Let $u \in W_{loc}^{1,2}(\mathbb{R}_+^{d+1})$ be a weak solution of $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$. Suppose that $N(\nabla u) \in L^p(\mathbb{R}^d)$ for some $1 < p < \infty$. Then u is a weak solution to the Neumann problem on $\mathbb{R}_+^{d+1}$ with data g for some $g \in L^p(\mathbb{R}^d)$. Furthermore, $\|g\|_p \leq C\|N(\nabla u)\|_p$ and*

$$-\frac{1}{\rho} \int_\rho^{2\rho} a_{(d+1)j}(x, t) \frac{\partial u}{\partial x_j} dt \rightarrow g \text{ weakly in } L^p(\mathbb{R}^d),$$

as $\rho \rightarrow 0^+$.

Proof. Let ψ be a Lipschitz function on $\mathbb{R}^d$ with compact support. Define

$$\Lambda(\psi) = \int_{\mathbb{R}_+^{d+1}} a_{ij} \frac{\partial u}{\partial x_j} \frac{\partial \varphi}{\partial x_i} dX, \quad (7.13)$$

where φ is a Lipschitz function on $\mathbb{R}^{d+1}$ with compact support such that $\varphi(x, 0) = \psi(x)$ for $x \in \mathbb{R}^d$. Since $N(\nabla u) \in L^p(\mathbb{R}^d)$, $|\nabla u| \in L^p(T(0, R))$ for any $R > 1$. Thus the integral in (7.13) converges. Also note that since $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$, the functional Λ is well defined.

Next let $\varphi(x, t) = \psi(x)\eta(t)$ in (7.13), where η is a continuous function such that $\eta(t) = 1$ on $[0, \rho]$, $\eta(t) = 0$ on $[2\rho, \infty)$ and η is linear on $[\rho, 2\rho]$. This gives

$$\Lambda(\psi) = \lim_{\rho \rightarrow 0} \left\{ -\frac{1}{\rho} \int_{\rho}^{2\rho} \int_{\mathbb{R}^d} \sum_{j=1}^{d+1} a_{(d+1)j} \frac{\partial u}{\partial x_j} \psi dx dt \right\}, \quad (7.14)$$

which, by Hölder inequality, implies that $|\Lambda(\psi)| \leq C \|N(\nabla u)\|_p \|\psi\|_{p'}$. Hence, there exists $g \in L^p(\mathbb{R}^d)$ such that $\|g\|_p \leq C \|N(\nabla u)\|_p$ and

$$\int_{\mathbb{R}_+^{d+1}} a_{ij} \frac{\partial u}{\partial x_j} \frac{\partial \varphi}{\partial x_i} dX = \Lambda(\psi) = \int_{\mathbb{R}^d} g \psi dx = \int_{\mathbb{R}^d} g \varphi dx.$$

□

8 L^2 estimates for the Neumann problem

To solve the L^2 Neumann problem, as in the cases of the Dirichlet and regularity problems, we need to impose a local solvability condition: for $x_0 \in \mathbb{R}^d$,

$$\int_{B(x_0, 1)} |N_1(\nabla u)|^2 dx \leq C_2 \left\{ \int_{B(x_0, 2)} |g|^2 dx + \int_0^2 \int_{B(x_0, 4)} |\nabla u|^2 dx dt \right\}, \quad (8.1)$$

whenever $u \in W^{1,2}(T(x_0, 4))$ satisfies (7.2) with $g \in C(B(x_0, 4))$ for all $\varphi \in C_0^1(B(x_0, 4) \times (-4, 4))$.

The goal of this section is to prove the following.

Theorem 8.1. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ with smooth coefficients satisfying (1.1), (1.2) and (1.3). Also assume that $\mathcal{L}$ satisfies conditions (4.1) and (8.1). Then, given any $g \in L^2(\mathbb{R}^d)$, there exists a $u \in W_{loc}^{1,2}(\mathbb{R}_+^{d+1})$, unique up to constants, satisfying $N(\nabla u) \in L^2(\mathbb{R}^d)$ and (7.2). Moreover the solution u satisfies the estimate $\|N(\nabla u)\|_2 \leq C \|g\|_2$, where C depends only on d , μ and the constants C_0 in (4.1) and C_2 in (8.1).*

In view of Theorems 7.4 and 7.7, it suffices to prove the following.

Theorem 8.2. *Under the same conditions on $\mathcal{L}$ as in Theorem 8.1, the L^2 Neumann problem for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable.*

Proof. Let $g \in C_c(\mathbb{R}^d)$ and u be given by (7.7). It follows from Lemma 7.2 that $u \in W^{1,2}(T(0, R))$ for any $R > 1$. This allows us to deduce from the condition (8.1) by an integration in x_0 over $\mathbb{R}^d$ that

$$\int_{\mathbb{R}^d} |N_4(\nabla u)|^2 dx \leq C \left\{ \int_{\mathbb{R}^d} |g|^2 dx + \int_0^6 \int_{\mathbb{R}^d} |\nabla u|^2 dx dt \right\}. \quad (8.2)$$

Note that by the pointwise estimates on $N(X, Y)$, we have $u(X) = O(|X|^{1-d})$ as $|X| \rightarrow \infty$. By Cacciopoli's inequality, this implies that

$$\iint_{T(0, 3R) \setminus T(0, R)} |\nabla u|^2 dx dt \rightarrow 0 \text{ as } R \rightarrow \infty.$$

In view of (3.6), we obtain

$$\int_0^6 \int_{\mathbb{R}^d} |\nabla u|^2 dx dt \leq C \int_{\mathbb{R}^d} |g|^2 dx. \quad (8.3)$$

By (8.2), this leads to $\|N_4(\nabla u)\|_2 \leq C\|g\|_2$.

Next, using the L^2 bound of $N_4(\nabla u)$ above and Lemma 5.4, we see that $u \rightarrow f$ n.t. on $\mathbb{R}^d$ and $\|\nabla_x f\|_2 \leq C\|N_4(\nabla u)\|_2 \leq C\|g\|_2$. Thus, by Remark 2.2,

$$\begin{aligned} \int_{\mathbb{R}^d} |N(\nabla u)|^2 dx &\leq C \left\{ \int_{\mathbb{R}^d} |\nabla_x u(x, 0)|^2 dx + \int_{\mathbb{R}^d} |N_4(\nabla u)|^2 dx + \int_{\mathbb{R}^d} |(Q(u))^*|^2 dx \right\} \\ &\leq C \left\{ \int_{\mathbb{R}^d} |g|^2 dx + \int_{\mathbb{R}^d} |(Q(u))^*|^2 dx \right\}. \end{aligned} \quad (8.4)$$

Finally we note that by (7.6), $|Q(u)(X)| \leq C|X|^{1-d-\alpha}$ for $|X|$ large. This implies that $(Q(u))^* \in L^2(\mathbb{R}^d)$. Since the L^2 Dirichlet problem for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable, we obtain

$$\int_{\mathbb{R}^d} |(Q(u))^*|^2 dx \leq C \int_{\mathbb{R}^d} |Q(u)|^2 dx \leq C \int_0^1 \int_{\mathbb{R}^d} |\nabla u(x, t)|^2 dx dt \leq C \int_{\mathbb{R}^d} |g|^2 dx$$

where we used (8.3) in the last step. In view of (8.4), we have proved that $\|N(\nabla u)\|_2 \leq C\|g\|_2$. $\square$

We end this section with a localization theorem.

Theorem 8.3. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ with coefficient matrix $A(X)$ satisfying (1.1)-(1.2). Suppose that $(N)_2$ and $(R)_2$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable. Let $u \in W^{1,2}(T(x_0, 6r))$ be a weak solution in $T(x_0, 6r)$ such that $N_r(\nabla u) \in L^2(B(x_0, 4r))$ and $\frac{\partial u}{\partial \nu} = g$ on $B(x_0, 4r)$. Then*

$$\int_{B(x_0, r)} |N_r(\nabla u)|^2 dx \leq C \int_{B(x_0, 3r)} |g|^2 dx + \frac{C}{r} \int_{T(x_0, 3r)} |\nabla u|^2 dX. \quad (8.5)$$

Proof. Let $\varphi \in C_0^\infty(B(x_0, 3r) \times (-3r, 3r))$ such that $\varphi = 1$ in $T(x_0, 2r)$ and $|\nabla\varphi| \leq C/r$. Let $w = u - \beta$ where β is the average of u over $T(x_0, 3r)$. Then for $X \in T(x_0, 2r)$,

$$\begin{aligned} w(X) &= \int_{\mathbb{R}_+^{d+1}} a_{ij} \frac{\partial}{\partial y_j} \{N(X, Y)\} \cdot \frac{\partial(w\varphi)}{\partial y_i} dY \\ &= \int_{\mathbb{R}^d} N(X, Y) g\varphi dY - \int_{\mathbb{R}_+^{d+1}} N(X, Y) a_{ij} \frac{\partial w}{\partial y_j} \cdot \frac{\partial\varphi}{\partial y_i} dY \\ &\quad + \int_{\mathbb{R}_+^{d+1}} a_{ij} \frac{\partial}{\partial y_j} \{N(X, Y)\} w \cdot \frac{\partial\varphi}{\partial y_i} dY \end{aligned} \quad (8.6)$$

With this representation formula, the rest of the proof similar to that of Theorem 6.10 in [21]. We omit the details. $\square$

Remark 8.4. Let $u \in W_{loc}^{1,2}(T(x_0, 6r))$ be a weak solution in $T(x_0, 6r)$ such that $N(\nabla u) \in L^2(B(x_0, 4r))$ and $\frac{\partial u}{\partial \nu} = 0$ on $B(x_0, 6r)$. Suppose that $(N)_2$ and $(R)_2$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ are solvable. Let E be the average of u over $T(x_0, 4r)$. It follows from Theorem 8.3 that

$$\begin{aligned} \left(\frac{1}{r^d} \int_{B(x_0, r)} |N_r(\nabla u)|^2 dx \right)^{1/2} &\leq C \left(\frac{1}{r^{d+1}} \int_{T(x_0, 3r)} |\nabla u|^2 dX \right)^{1/2} \\ &\leq \frac{C}{r} \left(\frac{1}{r^{d+1}} \int_{T(x_0, 3.1r)} |u - E|^2 dX \right)^{1/2} \leq \frac{C}{r^{d+2}} \int_{T(x_0, 3.2r)} |u - E| dX \\ &\leq \frac{C}{r^{d+1}} \int_{T(x_0, 3.2r)} |\nabla u| dX \leq \frac{C}{r^d} \int_{B(x_0, 4r)} |N(\nabla u)| dx, \end{aligned}$$

where we have used Cacciopoli's inequality, boundary regularity of weak solutions, and Poincaré inequality. This yields the weak reverse Hölder inequality,

$$\left\{ \frac{1}{r^d} \int_{B(x_0, r)} |N(\nabla u)|^2 dx \right\}^{1/2} \leq \frac{C}{r^d} \int_{B(x_0, 4r)} |N(\nabla u)| dx. \quad (8.7)$$

By the self-improving property of the weak reverse Hölder inequality, there exists $p > 2$, depending only on d and the constant C in (8.7), such that

$$\left\{ \frac{1}{r^d} \int_{B(x_0, r)} |N(\nabla u)|^p dx \right\}^{1/p} \leq C \left\{ \frac{1}{r^d} \int_{B(x_0, 2r)} |N(\nabla u)|^2 dx \right\}^{1/2}. \quad (8.8)$$

9 L^p estimates for the Neumann problem

In this section we study the L^p Neumann problem in $\mathbb{R}_+^{d+1}$.

Theorem 9.1. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ with coefficients satisfying (1.1)-(1.2). Suppose that $(N)_2$ and $(R)_2$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ are both solvable. Then $(N)_p$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable for $1 < p < 2 + \delta$.*

Proof. In the case of the unit ball (or a star-shaped Lipschitz domain), it was proved in [21] that the solvability of $(N)_{p_0}$ and $(R)_{p_0}$ implies the solvability of $(N)_p$ for any $1 < p < p_0 + \delta$. We follow the approach in [13] and [21] to treat the case $1 < p < 2$. For $f \in L^2(\mathbb{R}^d)$ with compact support, let $T(f) = N(\nabla u)$, where u is given by (7.7). Since $(N)_2$ is solvable, T is bounded on $L^2(\mathbb{R}^d)$. With Theorem 8.3 and pointwise estimates of the Neumann function at our disposal, one may establish $\|T(f)\|_1 \leq C$, where f is an H_{at}^1 atom. The L^p boundedness of T then follows by interpolation.

We use Theorem 6.7 to treat the range $p > 2$. Let $f \in L^2(\mathbb{R}^d)$ with compact support and $F = T(f) = N(\nabla u)$, where u is given by (7.7). For each ball B in $\mathbb{R}^d$, we choose $F_B = T(f\chi_{6B})$ and $R_B = T(f\chi_{\mathbb{R}^d \setminus 6B})$. Then $|F| \leq F_B + R_B$. By the L^2 boundedness of T ,

$$\int_B |F_B|^2 dx \leq C \int_{6B} |f|^2 dx.$$

To estimate R_B , we use the weak reverse Hölder inequality (8.8). This gives

$$\begin{aligned} \left(\frac{1}{r^d} \int_B |R_B|^p dx \right)^{1/p} &\leq C \left(\frac{1}{r^d} \int_{2B} |R_B|^2 dx \right)^{1/2} \\ &\leq C \left(\frac{1}{r^d} \int_{2B} |T(f)|^2 dx \right)^{1/2} + C \left(\frac{1}{r^d} \int_{6B} |f|^2 dx \right)^{1/2}. \end{aligned}$$

Thus, by Theorem 6.7, we obtain $\|T(f)\|_q \leq C\|f\|_q$ for any $2 < q < p$. □

The next theorem deals with the question of uniqueness for $1 < p < 2$.

Theorem 9.2. *Let $\mathcal{L}(u) = -\operatorname{div}(A\nabla u)$ with coefficients satisfying (1.1)-(1.2). Let $1 < p < 2$. Assume that $(R)_p$ and $(N)_2$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ are both solvable. Suppose that $N(\nabla u) \in L^p(\mathbb{R}^d)$ and $u \in W_{loc}^{1,2}(\mathbb{R}_+^{d+1})$ satisfies (7.11). Then u is constant in $\mathbb{R}_+^{d+1}$.*

Proof. Since $N(\nabla u) \in L^p(\mathbb{R}^d)$, $u = f$ n.t. on $\mathbb{R}^d$ for some $f \in \dot{W}^{1,p}(\mathbb{R}^d)$ by Lemma 5.4. Thus u is the solution of the L^p regularity problem with boundary data f . Choose $f_k \in C_0^1(\mathbb{R}^d)$ such that $\nabla_x f_k \rightarrow \nabla_x f$ in $L^p(\mathbb{R}^d)$. Let u_k be the solution of the classical Dirichlet problem with data f_k . By Theorem 7.8, u_k is also a weak solution to the Neumann problem on $\mathbb{R}_+^{d+1}$ with data $g_k \in L^p(\mathbb{R}^d)$. Since $\|g_k - g_\ell\|_p \leq C\|N(\nabla u_k - \nabla u_\ell)\|_p \rightarrow 0$ as $k, \ell \rightarrow \infty$, we may deduce that $g_k \rightarrow g$ in $L^p(\mathbb{R}^d)$.

Next we note that $\|N(\nabla u_k - \nabla u)\|_p \leq C\|\nabla_x(f_k - f)\|_p$. Thus $\nabla u_k \rightarrow \nabla u$ in $L^1(T(0, R))$ for any $R > 1$. It follows that $\int_{\mathbb{R}^d} g\varphi dx = 0$ for any $\varphi \in C_0^1(\mathbb{R}^{d+1})$. Consequently, we obtain $g = 0$.

Finally since $\|N(\nabla u_k)\|_2 \leq C\|\nabla_x f_k\|_2 < \infty$, u_k is a solution to the L^2 Neumann problem with data g_k . By Remark 7.5, u_k is given by $u_k = \beta_k + \int_{\mathbb{R}^d} N(X, y)g_k(y) dy$ for some constant β_k , if $d \geq 3$. If $d = 2$, one needs to replace $N(X, y)$ by $N(X, y) - N(X_0, y)$. In any case, because $g_k \rightarrow 0$ in $L^p(\mathbb{R}^d)$, we have $u_k(x) - \beta_k \rightarrow 0$ uniformly on any compact subset of $\mathbb{R}_+^{d+1}$. This, together with the fact that $\nabla u_k \rightarrow \nabla u$ in $L^1(T(0, R))$, implies that $\nabla u = 0$ a.e. in $\mathbb{R}_+^{d+1}$. Hence u is constant in $\mathbb{R}_+^{d+1}$. □

10 Proof of Theorems 1.1, 1.2 and 1.3

In this section we give the proof of Theorems 1.1, 1.2 and 1.3 stated in the Introduction. Let

$$D = \{(x, t) : x \in \mathbb{R}^d \text{ and } t > \psi(x)\},$$

where $\psi : \mathbb{R}^d \rightarrow \mathbb{R}$ is a Lipschitz function with Lipschitz constant $\|\nabla_x \psi\|_\infty$. Consider the bi-Lipschitzian mapping $\Phi : D \rightarrow \mathbb{R}_+^{d+1}$ given by $Y = \Phi(X) = \Phi(x, t) = (x, t - \psi(x))$. Let Φ' denote the Jacobian matrix of Φ and $\tilde{A}(Y) = (\tilde{a}_{ij}(Y)) = \Phi'(X)A(X)(\Phi'(X))^t$, where $X = \Phi^{-1}(Y) = \Psi(Y)$. Note that

$$\int_D a_{ij}(X) \frac{\partial u}{\partial x_j} \frac{\partial \varphi}{\partial x_i} dX = \int_{\mathbb{R}_+^{d+1}} \tilde{a}_{ij}(Y) \frac{\partial \tilde{u}}{\partial y_j} \frac{\partial \tilde{\varphi}}{\partial y_i} dY,$$

where $\tilde{u}(Y) = u(\Psi(Y))$ and $\tilde{\varphi}(Y) = \varphi(\Psi(Y))$. This shows that $\mathcal{L}(u) = -\operatorname{div}(A\nabla u) = 0$ in D if and only if $\tilde{\mathcal{L}}(\tilde{u}) = -\operatorname{div}(\tilde{A}\nabla \tilde{u}) = 0$ in $\mathbb{R}_+^{d+1}$. Consequently, the L^p Dirichlet problem for $\mathcal{L}(u) = 0$ in D is solvable if and only if the L^p Dirichlet problem for $\tilde{\mathcal{L}}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable. The same can be said for the L^p regularity and Neumann problems.

Note that $\Phi'(\Psi(Y))$ is independent of $s = y_{d+1}$ and $A(\Psi(Y)) = A(y, s + \psi(y))$. It follows that $\tilde{A}(y, s + 1) = \tilde{A}(y, s)$ if $A(x, t + 1) = A(x, t)$. Also, since

$$\tilde{A}(y, s_1) - \tilde{A}(y, s_2) = \Phi'(X) \{A(y, s_1 + \psi(y)) - A(y, s_2 + \psi(y))\} (\Phi'(X))^t,$$

there exist $c_1, c_2 > 0$ depending only on d and $\|\nabla_x \psi\|_\infty$ such that $c_1 \eta(\rho) \leq \tilde{\eta}(\rho) \leq c_2 \eta(\rho)$, where $\eta(\rho)$ is given by (1.10) and $\tilde{\eta}(\rho)$ is defined in the same manner, using $\tilde{A}$. Therefore the elliptic operator $\tilde{\mathcal{L}}$ satisfies the same assumptions as those imposed on $\mathcal{L}$ in Theorem 1.1. As a result, it suffices to prove Theorems 1.1, 1.2 and 1.3 in the case $D = \mathbb{R}_+^{d+1}$.

Next we reduce the general case to the case of smooth coefficients.

Lemma 10.1. *Let $\mathcal{L}_k = -\operatorname{div}(A_k \nabla)$ and $\mathcal{L} = -\operatorname{div}(A \nabla)$ with coefficient matrices satisfying (1.1)-(1.2). Suppose that $A_k(X) \rightarrow A(X)$ for a.e. $X \in \mathbb{R}^{d+1}$ as $k \rightarrow \infty$. Let u_k and u be solutions to the classical Dirichlet problem for $\mathcal{L}_k$ and $\mathcal{L}$ respectively in $\mathbb{R}_+^{d+1}$ with boundary data $f \in C_0^1(\mathbb{R}^d)$. Then, as $k \rightarrow \infty$, $u_k \rightarrow u$ uniformly in $T(0, R)$ for any $R > 1$ and $\nabla u_k \rightarrow \nabla u$ in $L^2(\mathbb{R}_+^{d+1})$. Consequently,*

$$\begin{aligned} \|(u)^*\|_p &\leq \liminf_{k \rightarrow \infty} \|(u_k)^*\|_p, \\ \|N(\nabla u)\|_p &\leq \liminf_{k \rightarrow \infty} \|N(\nabla u_k)\|_p. \end{aligned}$$

for any $p \geq 1$.

Proof. We may assume that $f \in C_0^1(\mathbb{R}^{d+1})$. Then

$$\int_{\mathbb{R}_+^{d+1}} A \nabla(u - f) \cdot \nabla g dX = - \int_{\mathbb{R}_+^{d+1}} A \nabla f \cdot \nabla g dX$$

for any $g \in C_0^1(\mathbb{R}_+^{d+1})$. Recall that $u(X) = \int_{\mathbb{R}^d} f d\omega^X = O(|X|^{1-d-\delta})$, as $|X| \rightarrow \infty$. By taking $g(X) = (u(X) - f(X))\varphi(X/R)$ where $\varphi \in C_0^\infty(\mathbb{R}^{d+1})$, $\varphi(X) = 1$ for $|X| < 1$ and $\varphi(X) = 0$ for $|X| \geq 2$, we may deduce that $\|\nabla u\|_{L^2(\mathbb{R}_+^{d+1})} \leq C\|\nabla f\|_{L^2(\mathbb{R}_+^{d+1})}$. Similarly, using

$$\int_{\mathbb{R}_+^{d+1}} A_k \nabla(u_k - u) \cdot \nabla g dX = - \int_{\mathbb{R}_+^{d+1}} (A_k - A) \nabla u \cdot \nabla g dX,$$

for any $g \in C_0^1(\mathbb{R}_+^{d+1})$, we obtain

$$\int_{\mathbb{R}_+^{d+1}} |\nabla(u_k - u)|^2 dX \leq C \int_{\mathbb{R}_+^{d+1}} |A_k - A|^2 |\nabla u|^2 dX.$$

It follows from the dominated convergence theorem that $\nabla u_k \rightarrow \nabla u$ in $L^2(\mathbb{R}_+^{d+1})$.

To see $u_k \rightarrow u$ uniformly in $T(0, R)$, we first note that $u_k \rightarrow u$ in $L^2(T(0, R))$. This follows from the Poincaré inequality

$$\int_{T(0, R)} |u_k - u|^2 dX \leq CR^2 \int_{T(0, R)} |\nabla(u_k - u)|^2 dX,$$

as $u_k - u = 0$ on $\mathbb{R}^d$. However, by De Giorgi -Nash estimates, the sequence $\{u_j\}$ is equi-continuous on $T(0, R)$. This implies that any subsequence of $\{u_k\}$ contains a subsequence which converges uniformly on $T(0, R)$ to u . It follows that $\{u_k\}$ converges uniformly on $T(0, R)$ to u . $\square$

Theorem 10.2. *Let $\mathcal{L}_k = -\operatorname{div}(A_k \nabla)$ and $\mathcal{L} = -\operatorname{div}(A \nabla)$ with coefficient matrices satisfying the same conditions as in Lemma 10.1. Suppose that $(R)_2$ for $\mathcal{L}_k(u_k) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable with uniform estimate $\|N(\nabla u_k)\|_2 \leq C_3 \|\nabla_x u_k\|_2$. Then $(R)_p$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable for $1 < p < p_0$, where $p_0 > 2$ depends only on d , μ and C_3 . Consequently, $(D)_q$ is solvable for $p'_0 < q < \infty$.*

Proof. By Theorem 6.1 and Theorem 5.7, we only need to show that $(R)_2$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable. To this end let u_k and u be the solutions to the classical Dirichlet problem for $\mathcal{L}(u) = 0$ and $\mathcal{L}_k(u_k) = 0$ in $\mathbb{R}_+^{d+1}$ with data $f \in C_0^1(\mathbb{R}^d)$, respectively. It follows from Lemma 10.1 and the uniform estimate $\|N(\nabla u_k)\|_2 \leq C_3 \|\nabla_x u_k\|_2$ that

$$\|N(\nabla u)\|_2 \leq \liminf_{k \rightarrow \infty} \|N(\nabla u_k)\|_2 \leq C_3 \|\nabla_x f\|_2.$$

$\square$

A similar result holds for the L^2 Neumann problem.

Lemma 10.3. *Let $\mathcal{L}_k = -\operatorname{div}(A_k \nabla)$ and $\mathcal{L} = -\operatorname{div}(A \nabla)$ with coefficient matrices satisfying the same conditions as in Lemma 10.1. Let*

$$\begin{aligned} u_k(X) &= \int_{\mathbb{R}^d} N_k(X, (y, 0)) f(y) dy, \\ u(X) &= \int_{\mathbb{R}^d} N(X, (y, 0)) f(y) dy, \end{aligned}$$

where $f \in L_c^\infty(\mathbb{R}^d)$ and $N_k(X, Y)$, $N(X, Y)$ are the Neumann functions constructed in Section 7 for $\mathcal{L}_k$, $\mathcal{L}$ respectively. Then $\nabla u_k \rightarrow \nabla u$ in $L^2(\mathbb{R}_+^{d+1})$ and consequently,

$$\|N(\nabla u)\|_p \leq \liminf_{j \rightarrow \infty} \|N(\nabla u_k)\|_p$$

for any $p \geq 1$.

Proof. The proof is similar to that of Lemma 10.1. Suppose that $\text{supp}(f) \subset B(0, r_0)$. Starting with

$$\int_{\mathbb{R}_+^{d+1}} A \nabla u \cdot \nabla g \, dX = \int_{\mathbb{R}^d} f g \, dx \quad \text{for } g \in C_0^\infty(\mathbb{R}^{d+1})$$

and using $u(X) = O(|X|^{1-d})$ as $|X| \rightarrow \infty$, we may deduce that

$$\int_{\mathbb{R}_+^{d+1}} |\nabla u|^2 \, dX \leq C r_0^{d+1} \|f\|_\infty^2.$$

Similarly, using

$$\int_{\mathbb{R}_+^{d+1}} A_k \nabla(u_k - u) \cdot \nabla g \, dX = - \int_{\mathbb{R}_+^{d+1}} (A_k - A) \nabla u \cdot \nabla g \, dX \quad \text{for } g \in C_0^\infty(\mathbb{R}^{d+1}),$$

we obtain

$$\int_{\mathbb{R}_+^{d+1}} |\nabla(u_k - u)|^2 \, dX \leq C \int_{\mathbb{R}_+^{d+1}} |A_k - A|^2 |\nabla u|^2 \, dX.$$

Thus $\nabla u_k \rightarrow \nabla u$ in $L^2(\mathbb{R}_+^{d+1})$ by the dominated convergence theorem. $\square$

Theorem 10.4. Let $\mathcal{L}_k = -\text{div}(A_k \nabla)$ and $\mathcal{L} = -\text{div}(A \nabla)$ with coefficient matrices satisfying the same conditions as in Lemma 10.1. Suppose that $(N)_2$ for $\mathcal{L}_k(u_k) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable with uniform estimate $\|N(\nabla u_k)\|_2 \leq C_4 \|\partial u_k / \partial \nu_k\|_2$. Then $(N)_2$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ is solvable. If in addition, $(R)_2$ is solvable for $\mathcal{L}_k(u_k) = 0$ with uniform estimate $\|N(\nabla u_k)\|_2 \leq C_4 \|\nabla_x u_k\|_2$, then both $(N)_p$ and $(R)_p$ for $\mathcal{L}(u) = 0$ in $\mathbb{R}_+^{d+1}$ are solvable for $1 < p < 2 + \delta$, where $\delta > 0$ depends only on n , μ and C_4 . Furthermore, the unique solutions of $(N)_p$ and $(R)_p$ satisfy $\|N(\nabla u)\|_p \leq C \|\partial u / \partial \nu\|_p$ and $\|N(\nabla u)\|_p \leq C \|\nabla_x u\|_p$ respectively with constant C depending only on d , p , μ and C_4 .

Proof. This follows readily from Lemma 10.3, Theorems 10.2 and 9.1. $\square$

Now given a $(d+1) \times (d+1)$ matrix $A(X)$ on $\mathbb{R}^{d+1}$ that satisfies conditions (1.1), (1.2), (1.3) and (1.11), we define

$$A_k(X) = \int_{\mathbb{R}^{d+1}} A(X - Y) \varphi_k(Y) \, dY$$

where $\{\varphi_k(Y)\}$ is a standard approximation of identity. Then $A_k \in C^\infty(\mathbb{R}^{d+1})$ and $A_k(X) \rightarrow A(X)$ for a.e. $X \in \mathbb{R}^{d+1}$. Furthermore, $A_k(X)$ satisfies the same conditions (1.1), (1.2), (1.3) and (1.11) with the same μ and $\eta_k(\rho) \leq \eta(\rho)$, where $\eta_k(\rho)$ denotes the modulus of continuity for A_k in the t variable. In view of Theorem 10.4, we have reduced the proofs of Theorems 1.1, 1.2 and 1.3 to the proof of the following theorem.

Theorem 10.5. *Theorems 1.1, 1.2 and 1.3 hold in the case where $D = \mathbb{R}_+^{d+1}$, $p = 2$, and the coefficient matrix $A(X)$ is C^∞ and satisfies conditions (1.1), (1.2), (1.3) and (1.11).*

Proof. In view of Theorems 4.1, 5.2 and 8.1, it suffices to show that the square Dini condition (1.11) implies the local solvability conditions (5.1) and (8.1). To this end, we fix $x_0 \in \mathbb{R}^d$ and choose $\varphi_1 \in C_0^\infty(\mathbb{R}^d)$ and $\varphi_2 \in C_0^\infty(\mathbb{R})$ so that $0 \leq \varphi_1 \leq 1$, $\varphi_1(x) = 1$ in $B(x_0, 8)$, $\varphi_1(x) = 0$ outside of $B(x_0, 9)$ and $0 \leq \varphi_2 \leq 1$, $\varphi_2(t) = 1$ in $[-8, 8]$, $\varphi_2(t) = 0$ outside of $[-9, 9]$. We consider the elliptic operator $L_1 = -\operatorname{div}(A_1 \nabla)$ in the star-like Lipschitz domain $\Omega_0 = T(x_0, 20)$, where

$$A_1(x, t) = \varphi_2(t) [\varphi_1(x) A(x, t) + (1 - \varphi_1(x)) I] + (1 - \varphi_2(t)) I \quad (10.1)$$

and I denotes the $(d+1) \times (d+1)$ identity matrix. Note that $A_1(x, t) = A(x, t)$ for $(x, t) \in T(x_0, 8)$ and thus $\mathcal{L}(u) = 0$ in $T(x_0, r)$ implies $L_1(u) = 0$ in $T(x_0, r)$, if $0 < r < 8$. We claim that under the condition (1.11), both $(N)_2$ and $(R)_2$ for $L_1(u) = 0$ in Ω_0 are solvable and the bounding constants in the nontangential maximal function estimates depend only on d , μ and $\eta(t)$ in (1.10). By the localization results for the L^p Neumann and regularity problems in star-like Lipschitz domains (see Theorems 5.19 and 6.10 in [21]), this gives estimates (5.1) and (8.1). We should point out that Theorems 5.19 and 6.10 in [21] were stated for solutions of $L_1(u) = 0$ in Ω_0 . However the same arguments in their proofs apply to local solutions of $L_1(u) = \mathcal{L}(u) = 0$ in $T(x_0, r)$ for $0 < r < 8$.

To prove the claim we introduce another elliptic operator $L_2 = -\operatorname{div}(A_2 \nabla)$ in Ω_0 , where

$$A_2(x, t) = \varphi_2(t) [\varphi_1(x) A(x, 0) + (1 - \varphi_1(x)) I] + (1 - \varphi_2(t)) I. \quad (10.2)$$

Let

$$\varepsilon^*(X) = \sup \{ |A_1(Y) - A_2(Y)| : Y \in B(X, \delta(X)/2) \}, \quad (10.3)$$

where $\delta(X) = \operatorname{dist}(X, \partial\Omega_0)$. Since

$$A_1(x, t) - A_2(x, t) = \varphi_1(x) \varphi_2(t) (A(x, t) - A(x, 0)), \quad (10.4)$$

it follows that $\varepsilon^*(X) = 0$ if $|X - P| \leq 1$ for some $P \in \partial\Omega_0 \setminus \{(y, 0) : |y| < 10\}$. It is also easy to see from (10.4) that if $X = (x, t)$ with $|x| \leq 11$ and $t \in (0, 1)$, then $\varepsilon^*(X) \leq \eta(2t)$, where $\eta(\cdot)$ is defined by (1.10). Therefore, for $P \in \partial\Omega_0$ and $r \in (0, 1)$,

$$h(P, r) := \frac{1}{r^d} \int_{B(P, r) \cap \Omega_0} \frac{(\varepsilon^*(X))^2}{\delta(X)} dX \leq C \int_0^{cr} \frac{(\eta(t))^2}{t} dt. \quad (10.5)$$

It follows from the condition (1.11) that

$$\sup \{ h(P, r) : P \in \partial\Omega_0 \} \rightarrow 0 \text{ as } r \rightarrow 0.$$

Hence, by Theorem 2.2 in [22], the solvability of $(N)_2$ and $(R)_2$ for $L_1(u) = 0$ in Ω_0 is equivalent to that of $(N)_2$ and $(R)_2$ for $L_2(u) = 0$ in Ω_0 . Thus it suffices to prove that the L^2 Neumann and regularity problems for $L_2(u) = 0$ in Ω_0 are solvable and the bounding constants in the nontangential maximal function estimates depend only on d and μ . We shall give the proof for the solvability of $(N)_2$. The proof for $(R)_2$ is similar.

Thus, let $u \in W^{1,2}(\Omega_0)$ be a weak solution to the Neumann problem for $L_2(u) = 0$ in Ω_0 with data $g \in C(\partial\Omega_0)$. We will show that

$$\int_{B(P,1) \cap \partial\Omega_0} |N(\nabla u)|^2 d\sigma \leq C \int_{\partial\Omega_0} |g|^2 d\sigma + C \int_{\Omega_0} |\nabla u|^2 dX \quad (10.6)$$

for any $P \in \partial\Omega_0$. The desired estimate $\|N(\nabla u)\|_2 \leq C\|g\|_2$ follows from (10.6) by a simple argument (see the proof of Theorem 1.5 in Section 11).

To see (10.6), we first note that $A_2(x, t) = I$ for $(x, t) \in \Omega_0 \setminus T(x_0, 9)$. Hence, if $P \in \partial\Omega_0 \setminus \{(y, 0) : |y| < 11\}$, u is harmonic in $B(P, 2) \cap \Omega_0$. As a result, estimate (10.6) follows readily from the solvability of the L^2 Neumann problem for $\Delta v = 0$ in Lipschitz domains [18] and Theorem 6.10 in [21]. To show (10.6) for $P = (y, 0)$ with $|y| < 11$, we use the fact that $A_2(x, t) = A_3(x, t)$ if $t \in [-8, 8]$, where

$$A_3(x, t) = \varphi_1(x)A(x, 0) + (1 - \varphi_1(x))I.$$

Let $L_3 = -\operatorname{div}(A_3 \nabla)$. Then $L_3(u) = 0$ in $B(x_0, 20) \times (0, 8)$. Since $A_3(X)$ is independent of the variable t , it follows from Theorem 8.1 in [21] (also see Theorem 1.12 in [1]) that both $(N)_2$ and $(R)_2$ for $L_3(v) = 0$ in $\mathbb{R}_+^{d+1}$ are solvable. Consequently, estimate (10.6) for $P = (y, 0)$ with $|y| \leq 11$ follows from Theorem 8.3. This concludes the proof of Theorem 10.5. $\square$

11 Uniform estimates in bounded Lipschitz domains

In this section we give the proof of Theorems 1.4, 1.5 and 1.6 stated in the Introduction. Let Ω be a bounded Lipschitz domain in $\mathbb{R}^{d+1}$, $d \geq 2$. Note that under the conditions (1.1), (1.2) and (1.14), the L^p Dirichlet problem for $\mathcal{L}_\varepsilon(u_\varepsilon) = 0$ is solvable in Ω for $2 - \delta < p < \infty$, while the L^p Neumann and regularity problems for $\mathcal{L}_\varepsilon(u_\varepsilon) = 0$ in Ω are solvable for $1 < p < 2 + \delta$. This follows directly from [12] and [21]. However, without the periodicity condition, the constants C (and δ) in the nontangential maximal function estimates in general depend on the parameter ε .

The proof of Theorems 1.4, 1.5 and 1.6 relies on the following two observations:

1. The class of elliptic operators $\{\mathcal{L}_1 = -\operatorname{div}(A \nabla)\}$ with $A(X)$ satisfying the conditions in Theorem 1.4 is invariant under rotation of the coordinate system.

2. Suppose $\mathcal{L}_\varepsilon(u_\varepsilon) = 0$ in $D = \{(x, t) : x \in \mathbb{R}^d \text{ and } t > \psi(x)\}$. Let $w(X) = u_\varepsilon(\varepsilon X)$. Then $\mathcal{L}_1(w) = 0$ in

$$D_\varepsilon = \{(x, t) \in \mathbb{R}^{d+1} : x \in \mathbb{R}^d \text{ and } t > \psi_\varepsilon(x)\},$$

where $\psi_\varepsilon(x) = \varepsilon^{-1}\psi(\varepsilon x)$.

Since $\|\nabla \psi_\varepsilon\|_\infty = \|\nabla \psi\|_\infty$, it follows from the observation (2) and Theorem 1.1 that solutions of the L^2 Dirichlet problem for $\mathcal{L}_\varepsilon(u_\varepsilon) = 0$ in D satisfies the estimate $\|(u_\varepsilon)^*\|_{L^2(\partial D)} \leq C\|u_\varepsilon\|_{L^2(\partial D)}$ uniformly in $\varepsilon > 0$. Similarly, by Theorems 1.2 and 1.3, solutions of the L^2 Neumann and regularity problems for $\mathcal{L}_\varepsilon(u_\varepsilon) = 0$ in D satisfy the uniform estimates

$$\|N(\nabla u_\varepsilon)\|_{L^2(\partial D)} \leq C\|\partial u_\varepsilon / \partial \nu_\varepsilon\|_{L^2(\partial D)} \quad \text{and} \quad \|N(\nabla u_\varepsilon)\|_{L^2(\partial D)} \leq C\|\nabla_{\tan} u_\varepsilon\|_{L^2(\partial D)},$$

respectively.

We begin with the Dirichlet problem for $\mathcal{L}_\varepsilon(u_\varepsilon) = 0$ in Ω .

Proof of Theorem 1.4. Although Theorem 1.6 implies Theorem 1.4, we give a direct proof.

Let ω_ε^X denote the $\mathcal{L}_\varepsilon$ -harmonic measure on $\partial\Omega$ and $K_\varepsilon(X, P) = d\omega_\varepsilon^X/d\sigma$, where $X \in \Omega$ and $P \in \partial\Omega$. Let $I(P, r) = B(P, r) \cap \partial\Omega$. We will show that for any $P_0 \in \partial\Omega$ and $0 < r < r_0$,

$$\left\{ \frac{1}{r^d} \int_{I(P_0, r)} |K_\varepsilon(X, P)|^2 d\sigma(P) \right\}^{1/2} \leq \frac{C}{r^d} \int_{I(P_0, r)} K_\varepsilon(X, P) d\sigma(P), \quad (11.1)$$

where $X \in \Omega \setminus B(P_0, 4r)$ and C is independent of ε .

Fix $P_0 \in \partial\Omega$ and $0 < r < r_0$. Since the class of elliptic operators $\{-\operatorname{div}(A\nabla)\}$ with $A(X)$ satisfying the conditions of Theorem 1.4 is invariant under rotation, we may assume that

$$\begin{aligned} \Omega \cap B(P_0, Cr_0) &= \{(x, t) : x \in \mathbb{R}^d \text{ and } t > \psi(x)\} \cap B(P_0, Cr_0) \\ &= D \cap B(P_0, Cr_0) \end{aligned} \quad (11.2)$$

and $A(x, t + t_0) = A(x, t)$ for some $t_0 \geq 1$ and any $(x, t) \in \mathbb{R}^{d+1}$. Without loss of generality, we may also assume that $t_0 = 1$. Let $G_\Omega(X, Y)$ and $G_D(X, Y)$ denote the Green's functions for $\mathcal{L}_\varepsilon$ on Ω and D respectively. Fix $Y \in D \setminus B(P_0, 4r)$. By the comparison principle (2.5), we have

$$\frac{G_\Omega(X, A_s(P))}{G_D(Y, A_s(P))} \approx \frac{G_\Omega(X, A_r(P_0))}{G_D(Y, A_r(P_0))} \quad \text{for } 0 < s < r \text{ and } P \in I(P_0, r),$$

where $A_s(P) = P + (0, \dots, 0, c_0 s)$. It follows that

$$\begin{aligned} \frac{\omega_\Omega^X(I(P, s))}{s^d} &\approx \frac{G_\Omega(X, A_s(P))}{s} \approx \frac{G_D(Y, A_s(P))}{s} \cdot \frac{G_\Omega(X, A_r(P_0))}{G_D(Y, A_r(P_0))} \\ &\approx \frac{\omega_D^Y(I(P, s))}{s^d} \cdot \frac{G_\Omega(X, A_r(P_0))}{G_D(Y, A_r(P_0))}, \end{aligned}$$

where ω_D denotes the $\mathcal{L}_\varepsilon$ -harmonic measure on ∂D . This implies that for any $P \in I(P_0, r)$,

$$K_\varepsilon(X, P) \leq C K_{\varepsilon, D}(Y, P) \cdot \frac{G_\Omega(X, A_r(P_0))}{G_D(Y, A_r(P_0))},$$

where $K_{\varepsilon, D}(Y, P) = d\omega_\varepsilon^Y/d\sigma$. Note that by Theorem 1.1 and a simple rescaling argument, $K_{\varepsilon, D}(Y, P)$ satisfies the reverse Hölder inequality with exponent 2 on $I(P_0, r)$. Thus,

$$\begin{aligned} \int_{I(P_0, r)} |K_\varepsilon(X, P)|^2 d\sigma(P) &\leq C \int_{I(P_0, r)} |K_{\varepsilon, D}(Y, P)|^2 d\sigma(P) \left(\frac{G_\Omega(X, A_r(P_0))}{G_D(Y, A_r(P_0))} \right)^2 \\ &\leq C r^{-d} [\omega_D^Y(I(P_0, r))]^2 \left(\frac{G_\Omega(X, A_r(P_0))}{G_D(Y, A_r(P_0))} \right)^2 \\ &\leq C r^{-d} [\omega_\Omega^X(I(P_0, r))]^2. \end{aligned}$$

This gives the desired reverse Hölder inequality (11.1). $\square$

Next we consider the L^p Neumann problem for $\mathcal{L}_\varepsilon(u_\varepsilon) = 0$ in Ω .

Proof of Theorem 1.5. By [21] it suffices to establish the uniform estimate $\|N(\nabla u_\varepsilon)\|_2 \leq C\|g\|_2$, where u_ε is the solution to the L^2 Neumann problem in Ω with boundary data $g \in L^2(\partial\Omega)$ and $\int_{\partial\Omega} g d\sigma = 0$. By dilation we may assume that $|\partial\Omega| = 1$.

Fix $P_0 \in \partial\Omega$. As in the proof of Theorem 1.4, we may assume that $\Omega \cap B(P_0, Cr_0)$ is given by (11.2) and $A(x, t+1) = A(x, t)$. By Theorem 1.2 and rescaling, the L^2 Neumann problem for $\mathcal{L}_\varepsilon(u) = 0$ in D is solvable with uniform estimates. It follows from Theorem 8.3 that

$$\int_{I(P_0, r_0)} |N(\nabla u_\varepsilon)|^2 d\sigma \leq C \int_{\partial\Omega} |g|^2 d\sigma + C \int_{\Omega} |\nabla u_\varepsilon|^2 dX.$$

By covering $\partial\Omega$ with balls of radius r_0 , we obtain

$$\int_{\partial\Omega} |N(\nabla u_\varepsilon)|^2 d\sigma \leq C \int_{\partial\Omega} |g|^2 d\sigma + C \int_{\Omega} |\nabla u_\varepsilon|^2 dX. \quad (11.3)$$

To estimate $\int_{\Omega} |\nabla u_\varepsilon|^2 dX$, we let β be the average of u_ε on Ω . Note that

$$\int_{\partial\Omega} |u_\varepsilon - \beta|^2 d\sigma \leq C \left\{ \int_{\Omega} |u_\varepsilon - \beta|^2 dX + \int_{\Omega} |\nabla u_\varepsilon|^2 dX \right\} \leq C \int_{\Omega} |\nabla u_\varepsilon|^2 dX,$$

where we have used the Poincaré inequality. Thus

$$\mu \int_{\partial\Omega} |\nabla u_\varepsilon|^2 d\sigma \leq \int_{\partial\Omega} g(u_\varepsilon - \beta) d\sigma \leq C(\rho) \int_{\partial\Omega} |g|^2 d\sigma + \rho \int_{\Omega} |\nabla u_\varepsilon|^2 dX,$$

where $0 < \rho < \mu$. This implies that $\|\nabla u_\varepsilon\|_{L^2(\Omega)} \leq C\|g\|_2$. In view of (11.3) we obtain $\|N(\nabla u_\varepsilon)\|_2 \leq C\|g\|_2$. $\square$

Finally we give the proof of Theorem 1.6.

Proof of Theorem 1.6. By [21] it suffices to show that

$$\|N(\nabla u_\varepsilon)\|_2 \leq C\|\nabla_{tan} f\|_2 + |\partial\Omega|^{\frac{1}{1-d}}\|f\|_2,$$

where u_ε is the solution of the L^2 regularity problem for $\mathcal{L}_\varepsilon(u_\varepsilon) = 0$ in Ω with data $f \in W^{1,2}(\partial\Omega)$. By dilation we may assume that $|\partial\Omega| = 1$.

By a localization argument similar to that in the proof of Theorem 1.5, we obtain

$$\int_{\partial\Omega} |N(\nabla u_\varepsilon)|^2 d\sigma \leq C \int_{\partial\Omega} |\nabla_{tan} f|^2 d\sigma + C \int_{\Omega} |\nabla u_\varepsilon|^2 dX.$$

Let $g = \frac{\partial u_\varepsilon}{\partial \nu_\varepsilon}$. The desired estimate follows from the observation

$$\mu \int_{\Omega} |\nabla u_\varepsilon|^2 dX \leq \rho\|g\|_2^2 + C(\rho)\|u_\varepsilon\|_2^2 \leq C\rho\|N(\nabla u_\varepsilon)\|_2^2 + C(\rho)\|f\|_2^2,$$

where we have used $\|g\|_2 \leq C\|N(\nabla u_\varepsilon)\|_2$. $\square$

Remark 11.1. Suppose $A(X)$ is periodic and $C^\alpha(\mathbb{R}^{d+1})$ for some $\alpha > 0$, Then it satisfies the condition (1.14). Furthermore, by the L^∞ gradient estimate in [2],

$$|\nabla u_\varepsilon(X)| \leq C \left(\frac{1}{|B(X, r)|} \int_{B(X, r)} |\nabla u_\varepsilon(Y)|^2 dY \right)^{1/2}$$

if $\mathcal{L}_\varepsilon(u_\varepsilon) = 0$ in $B(X, 2r)$. Consequently, the solutions of the L^p Neumann and regularity problems in Theorems 1.5 and 1.6 satisfy the uniform estimates

$$\|(\nabla u_\varepsilon)^*\|_p \leq C \|\partial u_\varepsilon / \partial \nu_\varepsilon\|_p \quad \text{and} \quad \|(\nabla u_\varepsilon)^*\|_p \leq C \|\nabla_{\tan} u_\varepsilon\|_p$$

for $1 < p < 2 + \delta$.

12 Appendix: Dahlberg's Theorem

In this appendix we give Dahlberg's original proof of Lemma 4.2 (and thus Theorem 4.1).

Lemma 12.1. *Let $\mathcal{L} = -\operatorname{div}(A\nabla)$ with coefficients satisfying (1.1), (1.2) and (1.3). Let u be a weak solution of $\mathcal{L}(u) = 0$ in $\Omega = B(x_0, r) \times (t_0 - r, t_0 + r)$ for some $(x_0, t_0) \in \mathbb{R}^{d+1}$ and $r > 3$. Then*

$$|u(x_0, t_0 + 1) - u(x_0, t_0)| \leq \frac{C}{r} \left(\frac{1}{|\Omega|} \iint_{\Omega} |u(y, s)|^2 dy ds \right)^{1/2}, \quad (12.1)$$

where C depends only on d and μ .

Proof. Let $F(x, t) = Q(u)(x, t)$. By the periodicity assumption (1.3), $\mathcal{L}(F) = 0$ in $B(x_0, r) \times (t_0 - r, t_0 + r - 1)$. It follows that

$$|F(x_0, t_0)| \leq C \left(\frac{1}{r^{d+1}} \int_{t_0 - \frac{r}{4}}^{t_0 + \frac{r}{4}} \int_{B(x_0, \frac{r}{4})} |F(y, s)|^2 dy ds \right)^{1/2}. \quad (12.2)$$

The desired estimate then follows from

$$|F(y, s)|^2 \leq \int_s^{s+1} \left| \frac{\partial u}{\partial t}(y, t) \right|^2 dt$$

and Cacciopoli's inequality. □

Lemma 12.2. *If in addition to the assumption in Lemma 12.1, we also assume that $u = 0$ in $\partial B(x_0, r) \times (t_0 - r, t_0 + r)$, then*

$$|u(x, t_0 + 1) - u(x, t_0)| \leq \frac{C}{r} \left(\frac{1}{|\Omega|} \iint_{\Omega} |u(y, s)|^2 dy ds \right)^{1/2} \quad (12.3)$$

for any $x \in B(x_0, r)$, where C depends only on d and μ .

Proof. The proof is similar to that of Lemma 12.1. In the place of (12.2), one uses the boundary L^∞ estimate. $\square$

Proof of Lemma 4.2.

The case $1 < r \leq 10$ follows easily by covering $B(x_0, r)$ with balls of radius 1. If $r > 10$, we also cover $B(x_0, r)$ with a sequence of balls $\{B(x_k, 1)\}$ of radius 1 such that

$$B(x_0, r) \subset \bigcup_k B(x_k, 1) \subset B(x_0, r+1) \quad \text{and} \quad \sum_k \chi_{B(x_k, 1)} \leq C.$$

Let I denote the left-hand side of (4.1). It follows from the assumption (4.1) (with $r = 1$) and boundary Harnack inequality (2.4) that

$$I \leq C \sum_k \int_{T(x_k, 2)} |u(X)|^2 dX \leq C \sum_k |u(x_k, 1)|^2. \quad (12.4)$$

To estimate the right-hand side of (12.4), we let v_1 be the Green's function for the operator $\mathcal{L}$ on the domain $\Omega_1 = B(x_0, 2r) \times (0, r)$ with pole at $(x_0, r/2)$. Similarly, we let v_2 be the Green's function for $\mathcal{L}$ on $\Omega_2 = B(x_0, 2r) \times (0, 2r)$ with pole at $(x_0, 3r/2)$. Note that

$$v_i(x_0, r/4) \geq cr^{2-d} \quad \text{for } i = 1, 2.$$

Thus, by the comparison principle (2.5), we have

$$\frac{u(x_k, 1)}{v_i(x_k, 1)} \leq C \frac{u(x_0, r/4)}{v_i(x_0, r/4)} \leq Cr^{d-2} u(x_0, r/4)$$

for $i = 1, 2$. In view of (12.4), this leads to

$$I \leq Cr^{2(d-2)} |u(x_0, r/4)|^2 \sum_k v_1(x_k, 1) v_2(x_k, 1). \quad (12.5)$$

Next we let ω denote the $\mathcal{L}$ -harmonic measure on the domain Ω_1 , evaluated at the point $(x_0, r/2)$. Since

$$\begin{aligned} v_1(x_k, 1) &\leq C \int_{B(x_k, 1) \times \{0\}} d\omega, \\ v_2(x_k, 1) &\leq C v_2(y, 1) \quad \text{for any } y \in B(x_k, 1), \end{aligned} \quad (12.6)$$

the right-hand side of (12.5) is bounded by

$$Cr^{2(d-2)} |u(x_0, r/4)|^2 \sum_k \int_{B(x_k, 1) \times \{0\}} v_2(y, 1) d\omega.$$

It follows that

$$I \leq Cr^{2(d-2)} |u(x_0, r/4)|^2 \int_{B(x_0, 2r) \times \{0\}} v_2(y, 1) d\omega. \quad (12.7)$$

We now consider the function

$$F(x, t) = v_2(x, t+1) - v_2(x, t) \quad \text{in } \Omega_1. \quad (12.8)$$

By the periodicity assumption (1.3), $\mathcal{L}(F) = 0$ in Ω_1 . It follows that

$$\begin{aligned} F(x_0, r/2) &= \int_{\partial\Omega_1} F d\omega \\ &= \int_{B(x_0, 2r) \times \{0\}} v_2(y, 1) d\omega + \int_{B(x_0, 2r) \times \{r\}} F(y, r) d\omega, \end{aligned} \quad (12.9)$$

where we have used the fact that $F = 0$ on $\partial B(x_0, 2r) \times (0, r)$ and $F(y, 0) = v_2(y, 1)$ on $B(x_0, 2r) \times \{0\}$. We claim that

$$|F(x_0, r/2)| \leq C r^{1-d}, \quad (12.10)$$

$$|F(y, r)| \leq C r^{1-d} \quad \text{if } |y - x_0| < 2r. \quad (12.11)$$

Assume the claim (12.10)-(12.11) for a moment. It then follows from (12.9) that

$$\int_{B(x_0, 2r) \times \{0\}} v_2(y, 1) d\omega \leq C r^{1-d}.$$

In view of (12.7), we obtain

$$I \leq C r^{d-3} |u(x_0, r/4)|^2 \leq \frac{C}{r^3} \int_{T(x_0, r)} |u(X)|^2 dX. \quad (12.12)$$

Finally we note that since $v_2(x, t) \leq C r^{2-d}$ for $(x, t) \in B(x_0, 2r) \times (0, r+3)$, estimates (12.10) and (12.11) follow readily from Lemmas 12.1 and 12.2, respectively. This completes the proof of Lemma 4.2. $\square$

References

- [1] M. A. Alfonseca, P. Auscher, A. Axelsson, S. Hofmann, and S. Kim, *Analyticity of layer potentials and L^2 solvability of boundary value problems for divergence form elliptic equations with complex L^∞ coefficients*, ArXiv:0705.0836v1.
- [2] M. Avellaneda and F. Lin, *Compactness methods in the theory of homogenization*, Comm. Pure Appl. Math. **40** (1987), 803–847.
- [3] ———, *Homogenization of elliptic problems with L^p boundary data*, Applied Math. Optim. **15** (1987), 93–107.
- [4] ———, *Compactness methods in the theory of homogenization II: Equations in nondivergent form*, Comm. Pure Appl. Math. **42** (1989), 139–172.
- [5] ———, *Homogenization of Poisson's kernel and applications to boundary control*, J. Math. Pure Appl. **68** (1989), 1–29.
- [6] ———, *L^p bounds on singular integrals in homogenization*, Comm. Pure Appl. Math. **44** (1991), 897–910.

- [7] A. Bensoussan, J.-L. Lions, and G.C. Papanicolaou, *Asymptotic Analysis and Periodic Structures*, North Holland, 1978.
- [8] L. Caffarelli, E. Fabes, S. Mortola, and S. Salsa, *Boundary behavior of non-negative solutions of elliptic operators in divergence form*, Indiana Univ. Math. J. **30** (1981), 621–640.
- [9] B. Dahlberg, personal communication.
- [10] ———, *On estimates of harmonic measure*, Arch. Ration. Mech. Anal. **65** (1977), 275–288.
- [11] ———, *On the Poisson integral for Lipschitz and C^1 -domains*, Studia Math. **66** (1979), 13–24.
- [12] ———, *On the absolute continuity of elliptic measure*, Amer. J. Math. **108** (1986), 1119–1138.
- [13] B. Dahlberg and C. Kenig, *Hardy spaces and the Neumann problem in L^p for Laplace's equation in Lipschitz domains*, Ann. of Math **125** (1987), 437–466.
- [14] R. Fefferman, *A criterion for the absolute continuity of the harmonic measure associated with an elliptic operator*, J. Amer. Math. Soc. **2** (1991), 127–135.
- [15] R. Fefferman, C. Kenig, and J. Pipher, *The theory of weights and the Dirichlet problem for elliptic equations*, Ann. of Math. **134** (1991), 65–124.
- [16] D. Jerison and C. Kenig, *An identity with applications to harmonic measure*, Bull. Amer. Math. Soc. **2** (1980), 447–451.
- [17] ———, *The Dirichlet problem in nonsmooth domains*, Ann. of Math. **113** (1981), 367–382.
- [18] ———, *The Neumann problem in Lipschitz domains*, Bull. Amer. Math. Soc. (N.S.) **4** (1981), 203–207.
- [19] C. Kenig, *Harmonic Analysis Techniques for Second Order Elliptic Boundary Value Problems*, CBMS Regional Conference Series in Math., vol. 83, AMS, Providence, RI, 1994.
- [20] C. Kenig, H. Koch, J. Pipher, and T. Toro, *A new approach to absolute continuity of elliptic measure, with applications to nonsymmetric equations*, Adv. Math. **153** (2000), 231–298.
- [21] C. Kenig and J. Pipher, *The Neumann problem for elliptic equations with non-smooth coefficients*, Invent. Math. **113** (1993), 447–509.
- [22] ———, *The Neumann problem for elliptic equations with nonsmooth coefficients: part II*, Duke Math. J. **81** (1995), 227–250.

- [23] C. Kenig and D. Rule, *The regularity and Neumann problem for non-symmetric elliptic operators*, Trans. Amer. Math. Soc. **361** (2009), 125–160.
- [24] C. Kenig and Z. Shen, *Boundary value problems for elliptic systems with periodic coefficients*, in preparation.
- [25] Z. Shen, *Necessary and sufficient conditions for the solvability of the L^p Dirichlet problem on Lipschitz domains*, Math. Ann. **336** (2006), no. 3, 697–724.
- [26] ———, *The L^p boundary value problems on Lipschitz domains*, Adv. Math. **216** (2007), 212–254.
- [27] ———, *A relationship between the Dirichlet and regularity problems for elliptic equations*, Math. Res. Letters **14** (2007), 205–213.
- [28] ———, *$W^{1,p}$ estimates for elliptic homogenization problems in nonsmooth domains*, Indiana Univ. Math. J. **57** (2008), 2283–2298.
- [29] G. Verchota, *Layer potentials and regularity for the Dirichlet problem for Laplace’s equation in Lipschitz domains*, J. Funct. Anal. **59** (1984), 572–611.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, IL 60637
E-mail address: cek@math.uchicago.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KENTUCKY, LEXINGTON, KY 40506
E-mail address: zshen2@email.uky.edu

August 14, 2009